



Penaeus aztecus Ives, 1891 (Crustacea, Decapoda), in the Scheldt estuary (Belgium): Isolated record or forerunner of a penaeid invasion?

Jan Soors^{a,*}, Jan Breine^a, Cédric d'Udekem d'Acoz^b, Ericia Van den Bergh^a, Frank Van de Meutter^a, Thomas Terrie^a

^a Research Institute for Nature and Forest (INBO), Herman Teirlinck Havenlaan 88 bus 73, 1000 Brussels, Belgium

^b Royal Belgian Institute of Natural Sciences, Service Heritage, Vautierstraat, 29, 1000 Brussels, Belgium

ARTICLE INFO

Keywords:

First record
Penaeus aztecus
Estuaries
Belgium
Invasive species

ABSTRACT

A single specimen of the penaeid prawn *Penaeus aztecus* (Ives, 1891) was recorded in 2018 in the brackish zone of the Scheldt estuary near Antwerp (Belgium). The presence of this species, native to the West Atlantic, might result either from ships' ballast water coming from transatlantic boat shipping, from illegal import or from a considerable expansion leap northwards from the Mediterranean Sea, where this species has recently established and now has rapidly expanding invasive populations.

1. Introduction

Invasions of marine and estuarine habitats by non-indigenous species are a worldwide phenomenon, which is steadily increasing as a consequence of human activities (Boudouresque, 2008; Galil et al., 2011; Ojaveer et al., 2018). The Scheldt estuary (The Netherlands - Belgium), with a high number of marine alien species recorded is no exception (e.g. Faasse and van Moorsel, 2003; Wolff, 2005; Kerckhof et al., 2007; Soors et al., 2010; De Blauwe, 2017). In the brackish zone we find established populations of over 50 non-indigenous species including many small crustaceans (Anonymous, 2010). The port of Antwerp (Belgium) renders this part of the river very vulnerable to introductions of non-indigenous species (Ysebaert et al., 1997). The main port activities of this large international harbor, receiving ships from over 800 locations worldwide, are situated in the brackish zone between the Dutch-Belgian border and the city of Antwerp.

The present paper reports the first record of a penaeid shrimp for the Scheldt estuary. Although the specimen was immature making identification difficult, all evidence indicates it to be a northern brown shrimp *Penaeus aztecus*. This species is native to the Western Atlantic (Pérez Farfante, 1969; Williams, 1984), has colonized the largest part of the Mediterranean Sea (Özcan et al., 2019), has recently reached the Black Sea (Gönülal and Türetken, 2019) and is now recorded for the first time in Northwest Europe.

2. Material and methods

The river Scheldt is 435 km long, originating on the plateau of Saint-Quentin in France and ending in the North Sea in the Netherlands near Vlissingen. The Scheldt estuary is approximately 160 km long and has a complete salinity gradient from polyhaline to a tidal freshwater zone, including extensive freshwater, brackish and saline tidal mudflats and marshes to its ecosystem. It is a well-mixed estuary characterized by strong currents, high turbidity and large tidal amplitude up to 6 m (Van den Bergh et al., 2005). Yearly, in the framework of the fish monitoring program by the Research Institute for Nature and Forest, fish assemblages are surveyed in spring, summer and fall with a mid-water beam trawl from an anchored boat. Each trawl consists of a net (mesh size 2 cm) fixed between two eight meter long steel beams. The lower beam is lowered to the bottom of the river while the upper beam is held at the surface. Both ends of the beams are attached to the anchor which keeps the boat at a fixed position. At the sampling site the nets are placed for two hours during flood as well as ebb-tide. Landed fish is measured and weighed and by-catch is counted and weighed. In Antwerp (Fig. 1) one single specimen of a penaeid shrimp was recorded during the 2018 fall survey. Its (tentative) identification is presented and the implications of the record are discussed.

2.1. Material examined

1 immature female (carapace length: 25 mm; total length 115 mm), Scheldt estuary, 65 km upstream, near the town of Antwerp,

* Corresponding author.

E-mail address: jan.soors@inbo.be (J. Soors).

<https://doi.org/10.1016/j.jembe.2020.151437>

Received 27 November 2019; Received in revised form 6 July 2020; Accepted 7 July 2020

0022-0981/ © 2020 Elsevier B.V. All rights reserved.




Fig. 1. Location of Belgian record of *Penaeus aztecus* (see arrow). *Penaeus aztecus* Ives, 1891.

51°12'19.84"N, 4°22'15.38"E (Fig. 1), mid-water beam trawl handled from an anchored boat, 27 Sept. 2018, deposited in the collections of the Royal Belgian Institute of Natural Sciences, registration number INV. 34094.

2.2. Characters and identification

The specimen was tentatively identified as *P. aztecus* based on Pérez Farfante (1969, 1970, 1988), Williams (1984), Tavares (2002), Deval et al. (2010) and Tavares and Gusmão (2016). It exhibited the following characters. Carapace smooth, rostrum armed with 10 dorsal teeth (epigastric spine included) and 2 ventral teeth; adrostral sulcus reaching almost to posterior margin of carapace; adrostral carina beginning slightly ahead of the first dorsal tooth on the rostrum and ends at 2 mm before of the hind edge of the carapace. Broad dorso-lateral sulcus on 6th abdominal somite. Telson without spines but with a deep dorsal sulcus (Fig. 2). The captured female measured 25 mm CL and 115 mm TL. Because this female was still juvenile—adult females *P. aztecus* measure up to 236 mm (TL)—, the identification of this one and only specimen is partly tentative and deductive. It definitely belongs to the species group previously referred as the genus *Farfantepenaeus*, and the developing thelycum has features in common with adult female *P. aztecus*: the posterior process is exposed and is armed with a median crest that bifurcates anteriorly to form an Y-shape (Pérez Farfante, 1969) (Fig. 3B). Additionally, the rostrum characters match with Pérez Farfante (1988): median sulcus is long and deep along the entire length and the dorsolateral sulcus is broad (Fig. 2A and B). The color of the freshly caught shrimp was not documented but the preserved specimen

was uniformly pale colored (Fig. 2), which suggests that it was a rather dull-colored shrimp, as reported for *P. aztecus* (Pérez Farfante, 1969). Additionally, considering the estuarine and temperate environment (*P. aztecus* is the only species of the *Farfantepenaeus* group occurring in cold-temperate waters (Williams, 1984)), the relative proximity of a population in the Mediterranean Sea and the invasive behavior of the species, the only likely species candidate is *Penaeus aztecus*. Despite all this, considering its immature status, an examination of its phenotype cannot confirm its identification with 100% certainty. The other highly unlikely candidates are species closely related to *P. aztecus* (other species of the former genus *Farfantepenaeus*), namely the northern pink shrimp *Penaeus duorarum* Burkenroad, 1939, *Penaeus isabelae* (Tavares and Gusmão, 2016) [no vernacular name], the southern pink shrimp *Penaeus notialis* Pérez Farfante, 1967, the southern brown shrimp *Penaeus subtilis* Pérez Farfante 1967 and the Sao Paulo shrimp *Penaeus paulensis* Pérez Farfante, 1967. These species were never found in Europe and have never been found far outside their home range.

2.3. Biology

Unlike most Penaeidae, *Penaeus aztecus* is not restricted to tropical and warm-temperate climatic conditions. In the West Atlantic, this species is present from the subtropical waters of Yucatan, Mexico and Belize up to cold temperate waters at Martha's Vineyard, Massachusetts, being the northern limit of this species (41°24'0"N) (Williams, 1984). Recently, it was recorded even further north at the Atlantic coasts of Canada.

It lives in coastal marine waters from the intertidal zone to



Fig. 2. A. Lateral anterior view of the Belgian *Penaeus aztecus*. B. Anterior dorsal view. C. Posterior dorsal view.

approximately 110 m deep, is rarely found at depths exceeding 165 m, and reaches its highest densities between 27 and 55 m, mainly on muddy, sandy-mud or loose peat bottoms (Pérez Farfante, 1969; Cook and Lindner, 1970; Williams, 1984). The species spawns offshore and the planktonic larvae and postlarvae are carried into estuaries by currents. They utilize estuarine marshes as nursery grounds. The juvenile shrimps grow within estuaries until they reach > 70 mm total length and then migrate offshore to mature and spawn (Knudsen et al., 1989; Matthews et al., 1991; Arreguin-Sanchez and Castro-Melendez, 2000). Larvae and juveniles are known to withstand a wide range of salinities and temperatures (Re et al., 2005).

3. Discussion

Among the growing list of alien species in the Belgian Scheldt estuary, *Palaemon macrodactylus* was the only invasive decapod shrimp or prawn that has been found so far (Soors et al., 2010).

Penaeus aztecus is the first penaeid prawn that has been found in Belgium. During the mid-water beam trawl of 27 September 2018, it was found among ten thousands of shrimps, mainly *Palaemon longirostris* (> 50.000 ind.) and *Crangon crangon* (> 20.000 ind.). The ichthyofauna on that day (20 species) was dominated by marine migrant fish such as the European sprat *Sprattus sprattus*, Atlantic herring *Clupea harengus*, European smelt *Osmerus eparlanus*, Sand goby *Pomatoschistus minutus* and Common goby *P. microps*. Exceptional were the capture of a Scaldfish *Arnoglossus laterna* (rare that far inland) and to a lesser extent the catches of Atlantic horse mackerel *Trachurus trachurus* and Sand smelt *Atherina boyeri* (Breine et al., 2019).

The northern brown shrimp is a commercially important Atlantic shrimp (Tavares, 2002). Globally 40.000 t was landed in 2010 (NOAA 2013). It has been introduced in New Caledonia and French Polynesia for aquacultural purposes (Scannella et al., 2017). Recently it was discovered in the Mediterranean Sea, where it has spread remarkably fast. From the Gulf of Antalya, where it first appeared, it has spread



Fig. 3. A. Lateral view of pleonite 6 and tail fan. B. Developing thelycum.

within eight years to the southeastern Levant, to the Gulf of Lion and to the Black Sea (Cruscanti et al., 2015; Galil et al., 2016; Zava et al., 2018; Gönülal and Türetken, 2019). It is suspected to compete with the native mediterranean penaeid prawn *Penaeus kerathurus* (Forskål 1775) (Kevrekidis, 2014). Both species reproduce in coastal waters, and post-larvae enter estuaries and coastal lagoons, where juveniles grow and later return to the marine environment (Scannella et al., 2017).

Eight of the fourteen non-indigenous penaeids in the Mediterranean were introduced through the Suez Canal (Galil et al., 2011) and they have become a new and increasingly important resource for mediterranean coastal fisheries (Scannella et al., 2017), owing to their high commercial value.

P. aztecus is frequently farmed in aquaculture facilities. Escape from shrimp farms or intentional release are probable introduction pathways. Other proposed vectors are ship's ballast water, illegal introduction and natural expansion by transportation of shrimp eggs and larvae along the main paths of surface water circulation and active swimming (Gönülal and Türetken, 2019). However, in the neighbouring countries, France (Le Havre) and the UK, the Kuruma prawn *Penaeus japonicus* (Spence Bate 1888) is the only *Penaeus* species that has been found with certainty on more than one occasion (Pezy et al., 2017). Another penaeid shrimp, *Penaeus semisulcatus* (De Haan 1844) was found only once, in August 2016 in Le Havre, English Channel (Pezy et al., 2017). These two species do not belong to the *Farfantepenaeus* group and are thus not to be confused with *P. aztecus*.

In the 2019 surveys no specimen of *P. aztecus* was caught. As the Scheldt estuary and surrounding water bodies are well monitored, it can be expected that the accidental or invasive nature of the presence of *P. aztecus* in the region will soon be established.

Author agreement

All authors have seen and approved the final version of the manuscript "Penaeus aztecus Ives, 1891 (Crustacea, Decapoda), in the Scheldt estuary (Belgium): isolated record or forerunner of a penaeid invasion?" that is submitted. We warrant that the article is the authors' original work, hasn't received prior publication and isn't under consideration for publication elsewhere.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors wish to thank the editors and reviewers for the constructive comments. We would like to thank the Flemish government for funding the mid-water beam trawl.

References

Anonymous, 2010. Bedreiging voor biodiversiteit. Indicatoren voor het Schelde-estuarium. Opgemaakt in opdracht van Afdeling Maritieme Toegang, projectgroep EcoWaMorSe, Vlaams Nederlandse Scheldecommissie. In: VLIZ Information Sheets, 200. Vlaams Instituut voor de Zee (VLIZ), Oostende, Belgium, pp. 7.

Arreguin-Sanchez, F., Castro-Melendez, R., 2000. On the interdependence of sequential fisheries: the brown shrimp, *Farfantepenaeus aztecus*, fisheries in the northwestern Gulf of Mexico. *Crustaceana* 73, 333–343. <https://doi.org/10.1163/156854000504426>.

Boudouresque, C.F., 2008. Les espèces introduites et invasives en milieu marin. Troisième édition. GIS Posidonie Publications, Marseille, pp. 201.

Breine, J., Galle, L., Lambeens, I., Maes, Y., Terrie, T., Van Thuyne, G., 2019. Opvolgen van het visbestand in het Zeeschelde-estuarium: Viscampagnes 2018. In: Rapporten van het Instituut voor Natuur- en Bosonderzoek. vol. 27 InstituutvoorNatuur-enBosonderzoek. <https://doi.org/10.21436/inbor.16173314>.

Cook, H.L., Lindner, M.J., 1970. Synopsis of biological data on the brown shrimp *Penaeus aztecus* Ives, 1891. *FAO Fish. Rep.* 57, 1471–1497.

Cruscanti, M., Innocenti, G., Bremer, J.A., Galil, B.S., 2015. First report of the brown shrimp *Penaeus aztecus* Ives, 1891 (Crustacea, Decapoda, Penaeidae) in the Tyrrhenian Sea. *Mar. Biodiver. Rec.* <https://doi.org/10.1017/S1755267215000664>.

De Blauwe, H., 2017. *Biftustra grandicella* (Canu and Bassler 1929) een warmteminnend mosdiertje (Bryozoa, Cheilostomata), nieuw voor Nederland en Europa. *De Strandvlo* 37 (1), 7–17.

Deval, M.C., Kaya, Y., Güven, O., Gökoglu, M., Frogli, C., 2010. An unexpected find of the Western Atlantic shrimp, *Farfantepenaeus aztecus* (Ives, 1891) (Decapoda, Penaeidae) in Antalya Bay, eastern Mediterranean Sea. *Crustaceana* 83 (12), 1531–1537. <https://doi.org/10.1163/001121610X538859>.

Faasse, M., van Moorsel, G., 2003. The North-American amphipods, *Melita nitida* Smith, 1873 and *Incisocalliope aestuarius* (Watling and Maurer 1973) (Crustacea: Amphipoda: Gammaridea), introduced to the Western Scheldt estuary (the Netherlands). *Aquat. Ecol.* 37, 13–22.

Galil, B.S., 2011. In: Galil, B., Clark, P., Carlton, J. (Eds.), Alien Crustaceans in the Mediterranean Sea: An Historical Review. In the Wrong Place - Alien Marine Crustaceans: Distribution, Biology and Impacts. Invading Nature - Springer Series in Invasion Ecology 6 Springer, Dordrecht. https://doi.org/10.1007/978-94-007-0591-3_13.

Galil, B.S., Innocenti, G., Douek, J., Paz, G., Rinkevich, B., 2016. Foul play? On the rapid spread of the brown shrimp *Penaeus aztecus* Ives, 1891 (Crustacea, Decapoda, Penaeidae) in the Mediterranean, with new records from the Gulf of Lion and the southern Levant. *Mar. Biodivers.* 47, 979–985. <https://doi.org/10.1007/s12526-016-0518-x>.

Gönülal, O., Türetken, P.S.Ç., 2019. One of the most invasive alien species, *Penaeus aztecus* Ives, 1891 reached the Black Sea coasts. *BioInvasions Rec.* 8 (4), 871–875. <https://doi.org/10.3391/bir.2019.8.4.15>.

Kerckhof, F., Haelters, J., Gollasch, S., 2007. Alien species in the marine and brackish ecosystem: the situation in Belgian waters. *Aquat. Invasions* 2 (3), 243–257. <https://doi.org/10.3391/ai.2007.2.3.9>.

Kevrekidis, K., 2014. The occurrence of the Atlantic penaeid prawn *Farfantepenaeus aztecus* (Ives, 1891) in the Thermaikos Gulf (Aegean Sea, Eastern Mediterranean): considerations on the potential establishment and impact on the autochthonous *Melicertus kerathurus* (Forskål, 1775). *Crustaceana* 87 (14), 1606–1619. <https://doi.org/10.1163/15685403-00003387>.

Knudsen, E.E., Paille, R.F., Rogers, B.D., Herke, W.H., Geaghan, J.P., 1989. Effects of a fixed-crest weir on brown shrimp *Penaeus aztecus* growth, mortality, and emigration in a Louisiana coastal marsh. *N. Am. J. Fish Manag.* 9, 411–419. [https://doi.org/10.1577/1548-8675\(1989\)009<0411:EOAFCW>2.3.CO;2](https://doi.org/10.1577/1548-8675(1989)009<0411:EOAFCW>2.3.CO;2).

Matthews, T.R., Schroeder, W.W., Stearns, D.E., 1991. Endogenous rhythm, light and salinity effects on postlarval brown shrimp *Penaeus aztecus* Ives recruitment to estuaries. *J. Exp. Mar. Biol. Ecol.* 154 (2), 177–189. [https://doi.org/10.1016/0022-0981\(91\)90163-Q](https://doi.org/10.1016/0022-0981(91)90163-Q).

NOAA, 2013. National Marine Fisheries Service. Monthly Commercial Landings Statistics Accessed 1 March 2020. http://www.st.nmfs.noaa.gov/st1/commercial/landings/monthly_landings.html.

Ojaveer, H., Galil, B.S., Carlton, J.T., Allevay, H., Gouletquer, P., Lehtiniemi, M., Marchini, A., Miller, W., Occhipinti-Ambrogi, A., Peharda, M., Ruiz, G.M., Williams, S.L., Zaiko, A., 2018. Historical baselines in marine bioinvasions: implications for policy and management. *PLoS One* 13 (8), e0202383. <https://doi.org/10.1371/journal.pone.0202383>.

Özcan, T., Ateş, A.S., Özcan, G., 2019. The distribution of the alien species *Penaeus aztecus* Ives, 1891 (Decapoda, Penaeidae) in the Mediterranean Sea. *Rev. Syst. Ecol. Res.* 21 (2), 41–48. <https://doi.org/10.2478/trser-2019-00011>.

Pérez Farfante, I., 1969. Western Atlantic shrimp of the genus *Penaeus*. U.S. fish and wildlife service. *Fish. Bull.* 67 (3), 461–591 i-x.

Pérez Farfante, I., 1970. Diagnostic characters of juveniles of the shrimps *Penaeus aztecus*, *P. duorarum duorarum*, and *P. brasiliensis* (Crustacea, Decapoda, Penaeidae). United States Fish and Wildlife Service. *Spec. Sci. Rep.* 599 (i–iii), 1–26.

Pérez Farfante, I., 1988. Illustrated key to Penaeoid shrimps of commerce in the Americas. *NOAA Tech. Rep.* 64 (i–iv), 1–32.

Pezy, P., Baffreau, A., Dauvin, J.-C., 2017. Records of two introduced Penaeidae (Crustacea: Decapoda) species from Le Havre Harbour. *BioInvasions Rec.* 6 (4), 363–367. <https://doi.org/10.3391/bir.2017.6.4.10>.

Re, A.D., Diaz, F., Sierra, E., Rodríguez, J., Perez, E., 2005. Effect of salinity and temperature on thermal tolerance of brown shrimp *Farfantepenaeus aztecus* (Ives) (Crustacea, Penaeidae). *J. Therm. Biol.* 30, 618–622. <https://doi.org/10.1016/j.jtherbio.2005.09.004>.

Scannella, D., Falsone, F., Geraci, M.L., Frogli, C., Fiorentino, F., Giusto, G.B., Zava, B., Insacco, G., Colloca, F., 2017. First report of northern brown shrimp *Penaeus aztecus* Ives, 1891 in strait of Sicily. *BioInvasions Rec.* 6 (1), 67–72. <https://doi.org/10.3391/bir.2017.6.1.11>.

Soors, J., Faasse, M., Stevens, M., Verbessem, I., De Regge, N., Van den Bergh, E., 2010. New crustacean invaders in the Schelde estuary (Belgium). *Belgian J. Zool.* 140 (1), 3–10.

Tavares, M., 2002. Shrimps. In: Carpenter, K.E. (Ed.), *The Living Resources of the Western Central Atlantic. Volume 1. Introduction, Molluscs, Crustaceans, Hagfishes, Sharks, Batoid Fishes and Chimaeras. FAO Species Identification Guide for Fishery Purposes – American Society of Ichthyologists and Herpetologists Special Publication 5. Food and agriculture Organization of the United Nations, Rome*, pp. 251–291.

Tavares, C., Gusmão, J., 2016. Description of a new Penaeidae (Decapoda: Dendrobranchiata) species, *Farfantepenaeus isabela* sp. nov. *Zootaxa* 4171 (3), 505–516. <https://doi.org/10.11646/zootaxa.4171.3.6>.

Van den Bergh, E., Van Damme, S., Graveland, J., de Jong, D., Baten, I., Meire, P., 2005. Ecological rehabilitation of the Schelde estuary (the Netherlands–Belgium; Northwest Europe): linking ecology, safety against floods, and accessibility for port

- development. *Restor. Ecol.* 13 (1), 204–214. <https://doi.org/10.1111/j.1526-100X.2005.00025.x>.
- Williams, A.B., 1984. *Shrimps, Lobsters, and Crabs of the Atlantic Coast of the Eastern United States, Maine to Florida.* i–xviii. Smithsonian Institution Press, Washington, D.C, pp. 1–550.
- Wolff, W.J., 2005. Non-indigenous marine and estuarine species in the Netherlands. *Zool. Mededelingen Leiden* 79, 1), 1–116.
- Ysebaert, T., Meire, P., De Block, M., De Regge, N., Soors, J., 1997. A first record of *Marenzelleria viridis* (Verril, 1873) (Polychaeta, Spionidae) in the Schelde estuary (Belgium). *Biol. Jaarboek (Dodonaea)* 64, 176–181.
- Zava, B., Insacco, G., Galil, B.S., 2018. The first record of the brown shrimp *Penaeus aztecus* Ives, 1891 in the central Adriatic coast of Italy. *Bioinvasions Rec.* 7 (3), 293–296. <https://doi.org/10.3391/bir.2018.7.3.10>.