

NEW LINGULIFORMEAN BRACHIOPODS FROM THE LOWER TREMADOCIAN (ORDOVICIAN) OF THE BRABANT MASSIF, BELGIUM, WITH COMMENTS ON CONTEMPORANEOUS FAUNAS FROM THE STAVELOT–VENN MASSIF

YVES CANDELA^{1*}, JEAN-MARC MARION², THOMAS SERVAIS³, WENHUI WANG⁴, MARK WOLVERS⁵ & BERNARD MOTTEQUIN⁶

¹*Corresponding author; Department of Natural Sciences, National Museums Scotland, Chambers Street, Edinburgh EH1 1JF, United Kingdom. E-mail: y.candela@nms.ac.uk

²Service géologique de Wallonie, Service public de Wallonie, Direction générale opérationnelle Agriculture, Ressources naturelles et Environnement, Avenue Prince de Liège, 15, B-5100 Jambes, Belgium. E-mail: jeanmarc.marion@spw.wallonie.be

³CNRS, Univ. Lille, UMR 8198 Evo-Eco-Paleo, F-59000 Lille, France. E-mail: thomas.servais@univ-lille.fr

⁴Key Laboratory of Metallogenic Prediction of Nonferrous Metals and Geological Environment Monitoring, Ministry of Education, School of Geosciences and Info-Physics, Central South University, Changsha 410083, PR China. E-mail: whwang@csu.edu.cn

⁵Brunel 19, 3641VG Mijdrecht, The Netherlands. E-mail: markwolvers@msn.com

⁶O.D. Earth and History of Life, Royal Belgian Institute of Natural Sciences, rue Vautier 29, B-1000 Brussels, Belgium.

E-mail: bmottequin@naturalsciences.be

To cite this article: Candela Y., Marion J.-M., Servais T., Wang W., Wolvers M. & Mottequin B. (2021) - New linguliformean brachiopods from the lower Tremadocian (Ordovician) of the Brabant Massif, Belgium, with comments on contemporaneous faunas from the Stavelot–Venn Massif. *Riv. It. Paleontol. Strat.*, 127(2): 383-395.

Keywords: Brachiopoda; Avalonia; Chevlipont Formation; low diversity; palaeogeography.

Abstract. Lower Ordovician brachiopod macrofaunas in Belgium (Avalonia) are seldom collected and studied due to the poor preservation of material. Here we describe a new fauna of linguliformean brachiopods from the Chevlipont Formation (lower Tremadocian) in the Brabant Massif. The fauna is of low diversity (at least three species belonging to *Rosobolus?*, *Thysanotos*, and *Broeggeria* have been identified) and is dominated by *B*. cf. *salteri* (Holl). Low diversity linguliformean brachiopod assemblages in a peri-Gondwanan terrane are characteristic of the lowermost Ordovician. Such assemblages are rooted in the Cambrian indicating that their geographic distribution during the early Ordovician was controlled by the radiation and dispersion of lineages surviving through the latest Cambrian–earliest Tremadocian linguliformean brachiopods taxonomic crisis. In addition we figure for the first time and comment on contemporaneous brachiopod faunas from the Stavelot-Venn Massif in SE Belgium. Finally, we present new graptolite data that enable a more precise constraint on the age for the Solwaster Member of the Jalhay Formation in the Stavelot-Venn Massif.

INTRODUCTION

Lower Palaeozoic rocks in Belgium crop out in six areas: the Brabant Massif, which constitutes the largest area (relative to its size it offers relatively less exposures) and occupies the central and

Received: February 8, 2021; accepted: April 28, 2021

north-western part of the country; the Condroz Inlier (also known as the Sambre–Meuse Strip) located in the centre; and the Ardennes in southern Belgium, with from the north-east to the south and south-east, the Stavelot–Venn, Serpont, Givonne and Rocroi massifs (also called inliers) (Fig. 1).

Contrary to the Belgian Devonian-Mississippian succession, the Lower Palaeozoic is poor in brachiopods, except for some Upper Ordovician



Fig. 1 - Location and schematic geological map of southern Belgium with indication of the localities cited in the text (modified from de Béthune (1954) and Mottequin (2021)). Abbreviations: Bi, Bierleux (Lhoest 1935; Vanguestaine et al. 2004); Ch, Chevlipont (this paper); G, Germany (see black and white insert); HSM OTS, Haine–Sambre–Meuse Overturned Thrust sheets (Belanger et al. 2012); LG, Lake Gileppe (e.g. Roncart 1925; Maillieux 1926; Schmidt & Geukens 1959); Li, Lierneux (Malaise 1878a, 1878b); N, the Netherlands (see black and white insert); So, Solwaster (e.g. Graulich 1963), Sp, Spa (Renier, unpubl.); Ta, Targnon (Charles 1925); TP, Trois-Ponts (Vanguestaine & Rushton 1979).

lithostratigraphic units (the Fosses Formation in the Condroz Inlier and the Huet Formation in the Brabant Massif) that has yielded diverse brachiopod faunas (e.g. Malaise 1873; Lespérance & Sheehan 1987; Sheehan 1987). Only one brachiopod occurrence was documented so far in the Cambrian (Wanne Formation) of the Stavelot–Venn Massif (Vanguestaine & Rushton 1979) (Fig. 2), with *Acrothele* cf. *bergeroni* Walcott, 1908 described from the Trois-Ponts locality (figured here on Fig. 3), whereas the Silurian, mostly represented by graptolitic shaly facies, is almost devoid of brachiopods (Maillieux 1930) although the Pridolian transgressive cover contains abundant but poorly diverse brachiopod faunas (e.g. Godefroid 1995; Mottequin 2019).

The lowermost Ordovician remained undiscovered in the Brabant Massif until the studies of Lecompte (1948, 1949), who recorded the presence of graptolitic horizons and scarce trilobite specimens in the Thyle Valley, demonstrating the presence of Tremadocian rocks. Since these studies no new macrofossil faunas have been described from the Lower Ordovician. Brachiopods are particularly rare in the Tremadocian of southern Belgium according to the literature. Several authors reported occurrences in the Jalhay Formation (Fig. 2) of the Stavelot–Venn Massif (Charles 1925; Roncart 1925; Mailllieux 1926) but these findings have never been illustrated until now. One of us (MW) has recently collected the material from the Chevlipont Formation we are describing below.

The bulk of the illustrated material is stored in the Royal Belgian Institute of Natural Sciences (Brussels; prefixed RBINS) and is complemented by specimens from the collections of the University of Liège (prefixed PA.ULg). Specimens selected for scanning electron microscopy were imaged with an ESEM FEI Quanta 200, under low vacuum; specimens were uncoated, except for *Acrothele* cf. *bergeroni*.



Fig. 2 - Chronostratigraphy and lithostratigraphy of the Brabant and Stavelot–Venn massifs (modified from Herbosch et al. 2020) with the distribution of the linguliformean brachiopods. The double arrows indicate boundaries between formations of uncertain age. Abbreviations: Brach., brachiopods; Gp., Groups; Syst., Systems.

GEOLOGY

Locality

The brachiopods from the Brabant Massif were collected by MW at the locality "Genappe 9" that is situated along the Charleroi-Ottignies railway, described by Lecompte (1949), in the Thyle Valley (Fig. 4). Lecompte (1949) collected a single specimen of Dictyonema flabelliforme var. sociale (Salter, 1858), re-identified as Rhabdinopora flabelliformis socialis (see Wang & Servais 2015). To date, no additional fauna was collected from locality "Genappe 9". This outcrop is located about 30 m north of Lecompte's locality "Genappe 78", located along what is now known as "Rue Basse Heuval", and about 30 m east of the site of the Moulin de Chevlipont. The locality "Genappe 78" has yielded numerous graptolites, more than 400 specimens according to Lecompte (1949: 5), who ascribed all of them to Dictyonema flabelliforme aff.

norvegica (Kjerulf, 1865), Wang & Servais (2015), however, identified a much more diverse assemblage with R. praeparabola Erdtmann, 1982, R. f. parabola (Bulman, 1954), R. f. flabelliformis (Eichwald, 1840), R. f. norvegica and R. f. socialis.

Stratigraphical summary

The locality investigated here lies within the Chevlipont Formation of the Ottignies Group (Fig. 2), in the Brabant Massif. Herbosch & Verniers (2013) and Herbosch & Debacker (2018) have provided detailed information on the geology of the Brabant Massif. The Chevlipont Formation (named by Anthoine & Anthoine 1943; see Verniers et al. 2002) consists typically of grey siltstone (also known as "quartzophyllade" in older literature) with millimetric to centimetric alternations of light grey siltstone and dark grey clayey siltstone and mudstone (Herbosch & Verniers 2013). This facies was interpreted as a low density tur-



Fig. 3 - Acrothele cf. bergeroni Walcott, 1908, PA.ULg 6464, Trois-Ponts (Stavelot–Venn Massif), Wanne Formation (Cambrian).

bidite typical of "mud turbidites" (see Herbosch et al. 2002); the Chevlipont Formation has been interpreted as a fine grained turbidite deposit on a depositional slope (Herbosch et al. 1991; André et al. 1991). The Chevlipont Formation shows a shallowing trend (Linnemann et al. 2012). Herbosch et al. (2002) also emphasised the known lack of benthic fauna in the formation and the presence of planktonic fauna (graptolites and acritarchs).

Work on dendroid graptolites and acritarchs (summary in Herbosch & Verniers 2013) has concluded an early Tremadocian age for the Chevlipont Formation. Wang & Servais (2015) have determined the lower boundary of the formation in the R. *praeparabola* graptolite Biozone, indicating an early Tremadocian age, Tr1 stage slice, mid 1a time slice (time slice of Webby et al. 2004) (for more details, see Herbosch & Verniers 2013).

Palaeogeographical implication

The Brabant Massif (Fig. 1) covers most of the northern part of Belgium and is unconformably covered by Devonian strata (Herbosch et al. 2002). It is within an orogenic zone called the Anglo-Brabant Deformation Belt, extending northwest under the North Sea into the concealed Caledonides of East Anglia (see Herbosch et al. 2002). During the Early Ordovician, the Brabant Massif was part of Avalonia, a peri-Gondwanan terrane located close to the northern margin of Gondwana (Amazonia-West Africa), at high latitudes. Murphy et al. (2006) proposed a rifting age for Avalonia and the opening of the Rheic Ocean constrained between the late Cambrian and the Early Ordovician. Shellnutt et al. (2019) correlated zircon populations with distinct magmatic periods in western Avalonia and more broadly across the whole of Avalonia. One of those magmatic 'pulses' correspond to rift-related magmatism (520 to 480 Ma) and the opening of the Rheic Ocean (Shellnutt et al. 2019). In addition, Herbosch & Verniers (2013) noted the decreasing sedimentation rate in lithostratigraphic units of the Brabant Massif during the late Cambrian and Early Ordovician (497 to 480 Ma) that announced the drifting of Avalonia from Gondwana (marked by the lower Tremadocian to Mid-Ordovician (Dapingian-Darriwilian) hiatus and unconformity) and the birth of the Rheic Ocean (Linnemann et al. 2012).

Early Ordovician macrofaunas are not commonly found in the Brabant Massif, as the thick lithostratigraphic succession (ranging from upper Terreneuvian to lower Tremadocian: 525–482 Ma) comprises clastic, mostly pelagic and turbiditic sediments deposited in an embayment of a large rift that developed on the western Gondwana margin (Linnemann et al. 2012) in continuation with the Pan-African orogen.

The brachiopod assemblage collected from the Chevlipont Formation is neither abundant nor diverse. Eighteen samples were collected totalling over 100 specimens, some of which cannot be fully identified due to their poor preservation. In decreasing abundance the assemblage is composed of *Broeggeria* cf. *salteri* (70% of the total amount of specimens identified), *Thysanotos* sp. (29%) and *Rosobolus?* sp. (1%). The brachiopods are not associated to graptolites and trilobites although these were reported by Lecompte (1949) in the investigated area.

The presence of this low diversity assemblage of linguliformean brachiopods in a peri-Gondwanan terrane, is typical of lowermost Ordovician brachiopod assemblages that have an origin in the Cambrian. Popov et al. (2013) suggested that the radiation and dispersion of lineages that survived the severe crisis linguliformean brachiopods experienced during the late Furongian to early Tremadocian clearly controlled the biogeographic distribution of this group in the Ordovician.

In the early Tremadocian, Thysanotos is only known from Belgium (present study). By the late Tremadocian to the earliest Floian, it is present in the South Urals and by the rest of the Floian, Thysanotos expanded its geographical distribution to northern Baltica (Estonia and Poland), the Armorica Terrane Assemblage (ATA) (see review of the concept in Servais & Sintubin 2013), and Iran; the last two being distributed along the northern margin of Gondwana. The Early Ordovician geographic extension of the genus is due to the palaeogeographical distribution of these terranes, with the southern part of Baltica and the South Urals in close vicinity of the northern margin of Gondwana and peri-Gondwanan terranes, at high southern latitudes, which may have helped larval expansion.

The genus *Broeggeria*, and in particular its type species *Broeggeria salteri*, is known from the Anglo-Welsh province of Avalonia during the late Cambrian–Tremadocian interval (Herefordshire, Merioneth Series; Wales, Tremadocian), and also contemporaneous horizons in the Kazakh terranes (Kendyktas Range, upper Cambrian–Tremadocian), Baltoscandia (Sweden, Norway, Denmark: upper Cambrian–upper Tremadocian), and Lau-



Fig. 4 - Detailed location map for the locality in the Chevlipont Formation (Brabant Massif).

rentia (Nova Scotia: Parabolina Zone, Furongian). The presence of this genus in both High-latitude and Low-latitude provinces, and in the Baltic Province (for the first two see sensu Harper et al. 2013) indicates a rapid geographic spread, or an older origin for this genus. By the Tremadocian, the geographical distribution of the genus has expanded to western peri-Gondwana (NW Argentina: B. omaguaca Benedetto et al., 2018; Santa Rosita Formation, see Benedetto et al. 2018), eastern peri-Gondwana (Bohemia: B. ferraria Mergl, 2002; Třenice Formation, see Mergl 2002), eastern Avalonia (Belgium: present study, Chevlipont and Jalhay formations) and the South Urals (B. salteri, lower part of the Kidryas Formation: upper Tremadocian-lower Floian).

Systematic Palaeontology

Subphylum LINGULIFORMEA

Williams et al., 1996 Class **LINGULATA** Gorjansky & Popov, 1985 Order **Lingulida** Waagen, 1885 Superfamily Linguloidea Menke, 1828 Family Zhanatellidae Koneva, 1986

Genus Rosobolus Havlíček, 1982 Type species: Rosobolus robertinus Havlíček, 1982, by original designation, from the Třenice Formation (Tremadocian), Holoubkov, Prague Basin, Czech Republic.

Rosobolus? sp.

Fig. 5.1

Material: One internal mould of a ventral valve.

Description. Ventral valve elongately oval with posterior margin angular; pseudointerarea small, apsacline. Surface of valve with fine, dense radial ornament, numbering 25 per mm. Interior with long, narrow pedicle groove, visceral area poorly preserved; central muscle scars not well defined, possibly subcircular or subrectangular, extending anteriorly for about 40% of valve length; *vascula lateralia* gently curved.

Remarks. The genera Rosobolus and Hyperobolus Havlíček, 1982 are similar but differ by the presence of large and subrectangular central muscle scars, roughly perpendicular to the midline of the ventral valve in the former, whereas in the latter these are shorter and oblique (see generic diagnosis of Rosobolus in Havlíček 1982: 19). Havlíček (1982) also emphasised the presence of fine radial capillae intersecting the short concentric lamellae ornamentation in Rosobolus only. Mergl (2002) highlighted the 'striking difference' in ornamentation between these two genera, with Rosobolus characterised by a smooth exterior, and a remarkably regular microornamentation, consisting of shallow transversely rhomboidal pits arranged in straight radial rows (these were first described by Holmer & Popov 2000: 64, fig. 29.1d). The tentative generic identification is based on the morphology of the central muscle scars and the presence of radial capillae. However, more material is needed to identify the genus with confidence.

Genus *Thysanotos* Mickwitz, 1896 Type species: *Obolus siluricus* von Eichwald, 1840, by original designation, from the Leetse Formation (Tremadocian–Floian), Paldiski, Estonia.

Thysanotos sp.

Fig. 5.2-6

Material: 44 internal and external moulds of both ventral and dorsal valves.

Description. Ventral valve suboval, 88% as long as wide, weakly convex; interior with visceral area subtriangular to slightly rhomboidal (anterior sides not as long as posterior sides); visceral area slightly raised anteriorly by a callosity, almost extending to mid valve length (Fig. 5.2); vascula lateralia divergent anteriorly, poorly preserved possibly extending forward to mid-valve length. Dorsal valve subcircular, 93% as long as wide, gently convex with maximum convexity towards the posterior third of valve length; interior not well-preserved, muscle scars hardly discernible; vascula lateralia widely divergent, only preserved in posterior half of the valve. Ornamentation poorly preserved, consisting of 9 concentric lamellae per mm, finer rugellae in the interspace (about 16 per mm); marginal spines infrequently preserved, and observed on one specimen (see Fig. 5.6).

Remarks. All the specimens collected from Belgium range from 5 to 7 mm in length; these are much smaller than specimens of *Thysanotos siluricus* in the type material of Bohemia (>30 mm), or specimens from Estonia (25-27 mm), Poland (20-22 mm) or the South Urals (40 mm). A second species from Bohemia, *Thysanotos primus* (Koliha, 1924), is smaller than *T. siluricus*, ranging from 16 to 19 mm long and never exceeding 20 mm (Mergl 1997).

The concentric ornament is also finer in the Belgian specimens (9 rugellae per mm) than in the samples from Bohemia (3-4 per mm for *T. primus*; 4-6 per mm for *T. siluricus*), north-eastern Europe (*T. siluricus* from Poland: 5-6 per mm; from Estonia: 4-6 per mm) and South Urals (4-6 per mm).

The distinction between *T. siluricus* and *T. primus* has focussed primarily on the difference in size and on the finer rugellate ornamentation on *siluricus* than on *primus*.



Fig. 5 - 1) Rosobolus? sp.; ventral valve interior, RBINS a13477. 2-6: Thysanotos sp.; 2) ventral valve interior, RBINS a13478; 3) dorsal valve interior RBINS a13479; 4) dorsal valve interior with impression of wavy concentric ornament, RBINS a13480; 5) dorsal valve exterior, RBINS a13481; 6) detail of external ornament with impression of marginal spines, RBINS a13482. 7-12: Broeggeria cf. salteri (Holl, 1865); 7) ventral valve interior, RBINS a13483; 8) ventral valve interior, RBINS a13484; 9) dorsal valve interior, RBINS a13486; 10) dorsal valve interior, RBINS a13486; 11) dorsal valve interior, RBINS a13487; 12 dorsal valve exterior, RBINS a13488. All the specimens are collected from the Chevlipont Formation, 2 km north of Villers-la-Ville, Brabant Massif.

Family Elkaniidae Walcott & Schuchert in Walcott, 1908

Genus Broeggeria Walcott, 1902

Type species: *Obolella salteri* Holl, 1865, by original designation, from the White-Leaved Oak Shale Formation (upper Cambrian), of the Malvern Hills, South Wales.

Broeggeria cf. salteri (Holl, 1865)

Fig. 5.7–12

cf. 1865 Obolella salteri Holl, p. 102, fig. 9.

Material: 89 internal and external moulds of both ventral and dorsal valves.

Description. Ventral valve subcircular, 85% as long as wide, gently convex in profile. Dorsal valve transversely suboval about 85% as long as wide, gently convex in lateral profile. Ornamentation consists of fine rugellae, numbering about 11-12 per mm. Dense faint radial striation is visible on interior moulds of many specimens (e.g. Fig. 5.7, 10, 11), but it is indistinguishable on the external moulds.

Ventral interior with short orthocline pseudointerarea; pedicle groove narrow, triangular, about 10% as wide as valve; visceral platform weakly thickened, rhomboidal in outline, with maximum width at anterior end, extending for about a third of valve length; *vascula lateralia* variably impressed in the present material, diverging anteriorly, flanking the visceral platform.

Dorsal interior characterised by a visceral area slightly raised above valve floor and extending beyond mid valve length; central muscle scars elongate (Fig. 5.9–11), gently divergent, bounded laterally by thin ridge, extending anteriorly for 45% of valve length; anterior lateral subcentral muscle scar, extending for 58% of valve length; *vascula media* subparallel to slightly divergent, not preserved anterior to visceral area; *vascula lateralia* widely divergent in posterior half of the valve, but anterior ends are not well preserved.

Remarks. *Monobolina* and *Broeggeria* have similar valve shapes and internal characters, but they differ externally in the presence of fila and costellae in *Monobolina* only (see diagnosis for *Monobolina* in Holmer & Popov 2000: 69). Although radial features are observed, in the present Belgian material, in internal moulds only (see also *Broeggeria salteri* in Popov & Holmer 1994: fig. 54D, F and 57D), these cannot be regarded as ornament as they are not expressed externally. These features are similar to the "drapes" described by Williams & Holmer (1992: 388), formed from stresses in the outer mantle groove induced by the spasmodic activity of "muscle sets controlling the setal follicles in the mantle grooves". Therefore, the specimens described here are best assigned to the genus *Broeggeria*.

The present specimens show a well-impressed dorsal visceral area, similar to that displayed in B. salteri, but this visceral area extends more anteriorly than in B. salteri from the Kendyktas Formation (upper Cambrian-Tremadocian) in the Kendyktas Range, Kazakhstan or from the Ceratopyge Shale (upper Tremadocian) in Öland, Sweden. Broeggeria salteri from the Tremadocian of South Wales (see Owens et al. 1982) and North Wales (see Sutton et al. 1999, 2000) are more transverse in outline than other populations of the same species and, in that feature, are more similar to the Belgian specimens. The present material is morphologically close to the type species of Broeggeria and thus here tentatively assigned to it. More specimens are nevertheless necessary to validate this comparison.

Comments on the Tremadocian Brachiopods from the Stavelot–Venn Massif

The first reports of linguliformean brachiopods within the Tremadocian of the Belgian part of the Stavelot-Venn Massif were published by Charles (1925) and Roncart (1925). Charles (1925) mentioned, without identifying them further than 'lingule', two specimens collected by students during an excursion. These specimens, which have not been traced in the collections of the Liège University, were collected from the locality with 'Dictyonema sociale' in the outcrop south of the confluent between the Lienne and Amblève rivers at Targnon (Fig. 1). Wang & Servais (2015) have re-assessed the graptolite faunas in the lower Tremadocian 'Dictyonema Shale' in Belgium and have re-identified 'Dictyonema sociale' as Rhabdinopora flabelliformis socialis, which in the Stavelot-Venn Massif, is found in the lower part of the Solwaster Member of the Jalhay Formation (for more information see Verniers et al. 2002) implying an early Tremadocian age (Tr1) for the occurrence (Fig. 2). However, Schmidt &



Fig. 6 - 1-2: *Broeggeria* sp.; 1) ventral valve interior in ventral (a) and oblique lateral (b) views, artificial cast of ventral interior in anteroventral (c) and slightly oblique anteroventral (d) views, and detail of visceral platform and *vascula lateralia* impression, RBINS a13489, Spa, route de Sart; 2). ventral valve exterior in ventral view (a), artificial cast of ventral valve exterior in ventral (b) and oblique lateral (c) views, PA.ULg 2020.12.16/1, Solwaster. 3) *Lingulella* sp., ventral valve interior in ventral (a) and oblique lateral (c, d) views and detail of pseudointerarea (b), RBINS a13490, Barrage de la Gileppe. 4) Pseudolingulidae indet., dorsal valve exterior in dorsal view (a) and detail of the ornamentation (b, c), PA.ULg 2020.12.16/2, Solwaster. All material from the Solwaster Member of the Jalhay Formation, Stavelot–Venn Massif. Specimen RBINS a13489 on Fig. 6.1a–b was coated with ammonium chloride; the rest of the material was imaged using a SEM (specimens uncoated).



Fig. 7 - 1-2: Graptolites associated with the linguliformean brachiopods from Solwaster (Stavelot–Venn Massif), Jalhay Formation (Solwaster Member). 1) ?Rhabdinopora praeparabola Erdtmann,1982, without proximal end but showing clear dissepiments (white arrows), PA.ULg 2020.12.16/3. 2) ?Rhabdinopora praeparabola Erdtmann,1982, a, juvenile form with clear planktonic sicula; b, specimen showing clearly divided nema/fibre structure, PA.ULg 2020.12.16/4.

Geukens (1959) critically commented on Charles' (1925) discovery, stressing that the graptolite he described was in fact Dictyonema flabelliforme flabelliforme. This form is also found in the lower part of the Solwaster Member of the Jalhay Formation (see Wang & Servais 2015). Roncart (1925) mentioned the discovery of another 'lingule' at the Lake Gileppe (Fig. 1); he identified it as the obolid Lingulella Salter, 1866, without precisely identifying the species (although specifying that it is close to L. cedens (Barrande)), due to its poor preservation (Fig. 6.3). Maillieux (1926) identified this obolid brachiopod from the "Assise de Vielsalm" (e.g. Graulich 1954) as Lingulella insons Barrande var. lata Koliha, confirming a Tremadocian age for this redundant stratigraphical unit that now corresponds to the Jalhay Formation (Verniers et al. 2002). Although they did not illustrate their material, Schmidt & Geukens (1959) reviewed and discussed the occurrence of Tremadocian brachiopods found on both sides of the Belgian-German border within the Stavelot-Venn Massif, i.e. in the Hürtgenwald region in Germany and around Lake Gileppe in Belgium. In summary, they recognised the presence of Lingulella insons, Acrotreta sp. in the strata with Rhabdinopora flabelliformis socialis, as well as Obolus (Bröggeria [sic]) salteri and an unidentified linguliformean brachiopod (mentioned as 'lingule') in the strata with Rhabdinopora flabelliformis flabelliformis. Both are from the uppermost lower Tremadocian (upper Tr1) (see Wang & Servais 2015: fig. 2 for the precise position of the horizon). Graulich (1963) reported Lingulella sp. and Obolus sp. at Solwaster (see also Geukens in Bulman 1970) (Fig. 1).

In the present study, we have identified more specimens from the Solwaster Member of the Jalhay Formation among the RBINS (Spa and Lake Gileppe; Fig. 1) and Liège University collections (Solwaster), namely Broeggeria sp. (Fig. 6.1-2), Lingulella sp. (Fig. 6.3) and Pseudolingulidae indet. (Fig. 6.4). Broeggeria from the Stavelot-Venn and Brabant massifs, the latter described and illustrated above (Fig. 5.7–12), possibly belong to the same species. However, more material from the Brabant Massif is needed to confirm this. The specimen identified as Pseudolingulidae indet. possesses similar external radial ornament to Meristopacha Sutton in Sutton et al., 1999, from the Anglo-Welsh Basin (Sutton et al. 1999) or Sedlecilingula Mergl, 1997 from Bohemia (Mergl 1997). However, the lack of more material does not permit a confident identification. Some graptolites were also found together with the brachiopod material from the Jalhay Formation at Solwaster (Fig. 1). These were identified by one of us (WW) as ?Rhabdinopora praeparabola (Fig. 7), indicating a mid early Tremadocian age (middle Tr1) for the assemblage (see Wang & Servais 2015).

The youngest Ordovician (early Floian) brachiopods from the Stavelot–Venn Massif are those reported by Malaise (1878a, 1878b) (?), Lhoest (1935) and Vanguestaine et al. (2004) from the Les Plattes Member of the Ottré Formation (Fig. 2) in the Lienne valley, at Bierleux and Lierneux (Fig. 1). Lhoest (1935) collected a single valve identified as *Lingulella* cf. *insons* by E. Maillieux (pers. comm. to Lhoest) (Les Plattes Member; see Vanguestaine et al.'s (2004) and Herbosch et al.'s (2020) discussions for the age of this level). Vanguestaine et al. (2004: pl. 1, fig. 17, 19-20) were the first to illustrate some of these poorly preserved linguliformean brachiopods.

Acknowledgements: The authors thank both reviewers, David Harper and Lars Holmer, for their comments that greatly improved the manuscript. We thank the editor, Lucia Angiolini, as well as Cristina Lombardo for dealing with the manuscript. The authors are grateful to Annelise Folie (RBINS) for providing the accession numbers for the specimens, Julien Cillis (RBINS) for the SEM photographic work, and to Valentin Fischer and Julien Denayer for access to the palaeontological collections of the University of Liège. YC thanks Eoghan Candela for drafting Figure 4.

This paper is a contribution to IGCP project 653 "The onset of the Great Ordovician Biodiversification Event", and to IGCP project 735 "Rocks and the Rise of Ordovician Life: Filling knowledge gaps in the Early Palaeozoic Biodiversification".

References

- André L., Herbosch A., Louwye S., Servais T., Van Grootel G., Vanguestaine M. & Verniers J. (1991) - Guidebook to the excursion on the stratigraphy and magmatic rocks of the Brabant Massif, Belgium. *Annales de la Société géologique de Belgique*, 114(2): 283-323.
- Anthoine R. & Anthoine P. (1943) Les assises de Mousty et de Villers-la-Ville du bassin supérieur de la Dyle. Annales de la Société géologique de Belgique, 66: M53-M181.
- Belanger I., Delaby S., Delcambre B., Ghysel P., Hennebert M., Laloux M., Marion J.-M., Mottequin B. & Pingot J.-L. (2012) - Redéfinition des unités structurales du front varisque utilisées dans le cadre de la nouvelle Carte géologique de Wallonie (Belgique). *Geologica Belgica*, 15(3): 169-175.
- Benedetto J. L., Lavié F.J. & Muñoz D.F. (2018) Broeggeria Walcott and other upper Cambrian and Tremadocian linguloid brachiopods from NW Argentina. Geological Journal, 53: 102-119.
- de Béthune P. (1954) Carte géologique de Belgique (échelle 1/500.000). Atlas de Belgique, planche 8. Académie royale de Belgique, Bruxelles.
- Bulman O.M. (1954) The graptolite fauna of the *Dictyonema* Shales of the Oslo region. *Norsk Geologisk Tidsskrift*, 33: 1-40.
- Bulman O.M. (1970) A new Dictyonema fauna from the Salmien [sic] of the Stavelot Massif. Bulletin de la Société belge de Géologie, de Paléontologie et d'Hydrologie, 79(3/4): 213-224.
- Charles F. (1925) Découverte de lingule dans le Salmien de la Lienne. *Annales de la Société géologique de Belgique*, 47: B197-198.
- von Eichwald C.E. (1840) O siluriyskoy systeme plastov v Estlyandii (pererabotannyy perevod s nemetskogo) [Ueber das silurische Schichtensystem in Esthland]., St. Petersburg. 210 pp.
- Erdtmann B.D. (1982) A reorganization and proposed phylogenetic classification of planktic Tremadoc (early Or-

dovician) dendroid graptolites. Norsk Geologisk Tidsskrift, 62: 121-145.

- Godefroid J. (1995) Dayia shirleyi Alvarez & Rachebœuf, 1986, un brachiopode silurien dans les 'Schistes de Mondrepuis' à Muno (sud de la Belgique). Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre, 65: 269-272.
- Gorjansky V.Iu. & Popov L.E. (1985) Morfologiia, systematicheskoe polozhenie I proiskhozhdenie bezzamkovykh brakhiopods karbonatnoi rakovinoi [Morphology, systematic position and origin of the inarticulate brachiopods with calcareous shells]. *Paleontologicheskii Zhurnal*, 1985(3): 3-14 [in Russian].
- Graulich J.M. (1954) Le Cambrien-Trémadocien. In: Fourmarier P. (Ed.) - Prodrome d'une description géologique de la Belgique : 21-38. Société géologique de Belgique, Liège.
- Graulich J.M. (1963) Découverte du niveau à *Dictyonema flabelliforme parabola* dans le Salmien du massif de Stavelot (Belgique). *Compte rendus hebdomadaires des séances de l'Académie des sciences*, 256: 3327-3328.
- Harper D.A.T., Rasmussen C.M.Ø., Liljeroth M., Blodgett R.B., Candela Y., Jin Jisuo, Percival I.G., Rong Jiayu, Villas E. & Zhan, Renbin (2013) - Biodiversity, biogeography and phylogeography of Ordovician rhynchonelliform brachiopods. In: Harper D.A.T. & Servais, T. (Eds)
 - Early Palaeozoic Biogeography and Palaeogeography. *Geological Society, London, Memoirs*, 38: 127-144.
- Havlíček V. (1982) Lingulacea, Paterinacea, and Siphonotretacea (Brachiopoda) in the Lower Ordovician sequence of Bohemia. Sborník geologických věd, Paleontologie, 25: 9-82.
- Herbosch A. & Debacker T.N. (2018) A new geological map of the outcrop areas of the Brabant Massif (Belgium). *Geologica Belgica*, 21(1/2): 41-58.
- Herbosch A., Liégeois J.-P., Gärtner A., Hofmann M. & Linnemann U. (2020) - The Stavelot–Venn Massif (Ardenne, Belgium), a rift shoulder basin ripped off the West African craton: Cartography, stratigraphy, sedimentology, new U–Pb on zircon ages, geochemistry and Nd isotopes evidence. *Earth-Science Reviews*, 203. https:// doi.org/10.1016/j.earscirev.2020.103142
- Herbosch A., Vanguestaine M., Degardin J. M., Dejonghe L., Fagel N. & Servais T. (1991) - Etude lithostratigraphique, biostratigraphique et sédimentologique du sondage de Lessines (bord meridional du Massif du Brabant, Belgique). Annales de la Société géologique de Belgique, 114(1): 195-212.
- Herbosch A. & Verniers J. (2013). Stratigraphy of the Lower Palaeozoic of the Brabant Massif, Belgium. Part 1: the Cambro-Ordovician from the Halle and Ottignies groups. *Geologica Belgica*, 16(1/2): 49-65.
- Herbosch A., Verniers J., Debacker T., Billiaert B., De Schepper S & Belmans M. (2002) - The Lower Palaeozoic stratigraphy and sedimentology of the Brabant Massif in the Dyle and Orneau Valleys and of the Condroz Inlier at Fosses: an excursion guidebook. *Geologica Belgica*, 5(3/4): 71-143.
- Holl H.B. (1865) On the geological structure of the Malvern

Hills and adjacent districts. *Quarterly Journal of the Geological Society, London,* 21: 72-102.

- Holmer L.E. & Popov L.E. (2000) Lingulida. In: Kaesler, R. L. (Ed.) - Treatise on Invertebrate Paleontology, Part H, Brachiopoda, Revised, Volume 2: Linguliformea, Craniiformea, and Rhynchonelliformea (part): 32-97. The Geological Society of America and The University of Kansas. Boulder, Colorado and Lawrence, Kansas.
- Kjerulf T. (1865) Veiviser ved geologiske excursioner i Christiana omegn. Universitets program for andet halvaar: 1-43. Brøgger & Christie's Bogtrykkeri, Christiania.
- Koliha J. (1924) Atremata z kruišnohorských vrstev (dα). *Palaeontographica Bohemiae*, 10: 1-61.
- Koneva S.P. (1986) Novoe semeistvo kembriiskikh bezzamkovykh brakhiopod [A new family of the Cambrian inarticulate brachiopods]. *Paleontologicheskii Zhurnal*, 1986(1): 49-55 [in Russian].
- Lecompte M. (1948) Existence du Trémadocien dans le Massif du Brabant. *Bulletin de l'Académie royale de Belgique, Classe des Sciences*, 34, 5e série: 677-687.
- Lecompte M. (1949) Découverte de nouveaux gîtes à *Dictyonema* dans le Trémadocien du Massif du Brabant. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 25(45): 1-8.
- Lespérance P.J. & Sheehan P.M. (1987) Trilobites et Brachiopodes ashgilliens (Ordovicien supérieur) de l'« Assise » de Fosse, Bande de Sambre-Meuse (Belgique). *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre*, 57: 91-123.
- Linnemann U., Herbosch A., Liégeois J.-P., Pin C., Gärtner A. & Hofmann M. (2012) - The Cambrian to Devonian odyssey of the Brabant Massif within Avalonia: A review with new zircon ages, geochemistry, Sm-Nd isotopes, stratigraphy and palaeogeography. *Earth-Science Reviews*, 112(2012): 126-154.
- Lhoest A. (1935) Sur la présence de « *Lingulella* cf. *insons* » dans le Salmien supérieur et nouveaux gîtes à « *Dictyonema flabelliforme* » dans la vallée de la Lienne. *Annales de la Société géologique de Belgique*, 58: B151-152.
- Maillieux E. (1926) Remarques sur l'Ordovicien de la Belgique. Bulletin de la Société belge de Géologie, de Paléontologie et d'Hydrologie, 36, Procès-Verbaux: 67-85.
- Maillieux E. (1930) Observations nouvelles sur le Silurien de Belgique. *Bulletin du Musée royal d'Histoire naturelle de Belgique*, 6(15): 1-8.
- Malaise C. (1873) Description du terrain silurien du centre de la Belgique. Mémoires Couronnés et Mémoires des Savants Étrangers, publiés par l'Académie royale des Sciences, des Lettres et des Beaux-Arts de Belgique, 37: 1-122.
- Malaise C. (1878a) Découverte de Brachiopodes du genre Lingula dans le cambrien [sic] du massif de Stavelot. Bulletin de l'Académie royale des Sciences, des Lettres et des Beaux-Arts de Belgique, 46, 2e série: 58.
- Malaise C. (1878b) Sur des Lingula trouvées à Lierneux, dans le cambrien [sic] de l'Ardenne. Annales de la Société géologique de Belgique, 5, Bulletin: 137-138.
- Menke C.T. (1828) Synopsis methodica molluscorum generum omnium et specierum earum quae in Museo Menkeano adservantur. G. Uslar, Pyrmonti, 91 pp.

- Mergl M. (1997) Distribution of the lingulate brachiopod *Thysanotos* in Central Europe. *Věstník Ceského geologického* ústavu, 72(1): 27-35.
- Mergl M. (2002) Linguliformean and craniiformean brachiopods of the Ordovician (Třenice to Dobrotivá formations) of the Barrandian, Bohemia. Acta Musei Nationalis Pragae, Series B, Natural History, 58(1/2): 1-82.
- Mickwitz A. (1896) Über die Brachiopodengattung Obolus Eichwald. Mémoires de l'Académie Impériale des sciences de St. Petersbourg, 4(2): 1-275.
- Mottequin B. (2019) An annotated catalogue of types of Silurian–Devonian brachiopod species from southern Belgium and northern France in the Royal Belgian Institute of Natural Sciences (1870–1945), with notes on those curated in other Belgian and foreign institutions. *Geologica Belgica*, 22(1/2): 47-89.
- Mottequin B. (2021) Earth science collections of the Centre Grégoire Fournier (Maredsous) with comments on Middle Devonian–Carboniferous brachiopods and trilobites from southern Belgium. *Geologica Belgica*, 24(1/2): 33-68.
- Murphy J.B., Gutierrez-Alonso G., Nance R.D., Fernandez-Suarez J., Duncan Keppie J., Quesada C., Strachan R.A.
 & Dostal J. (2006) Origin of the Rheic Ocean: Rifting along a Neoproterozoic suture? *Geology*, 34(5): 325-328.
- Owens R.M, Fortey R.A., Cope J.C.W., Rushton A.W.A. & Bassett M.G. (1982) Tremadoc faunas from the Carmarthen district, South Wales. *Geological Magazine*, 119(1): 1-38.
- Popov L.E. & Holmer L.E. (1994) Cambrian–Ordovician lingulate brachiopods from Scandinavia, Kazakhstan, and South Ural Mountains. *Fossils & Strata*, 35: 1-156.
- Popov L.E., Holmer L.E., Bassett M.G., Ghobadi Pour M. & Percival I.G. (2013) - Biogeography of Ordovician linguliform and craniiform brachiopods. In: Harper D.A.T. & Servais T. (Eds) - Early Palaeozoic Biogeography and Palaeogeography. *Geological Society, London, Memoirs,* 38: 117-126.
- Roncart Ch. (1925) Observation. *Annales de la Société géologique de Belgique*, 47: B198.
- Salter J.W. (1858) On Graptopora, a new genus of Polyzoa, allied to the graptolites. Proceedings of the American Association for the Advancement of Science (for 1857), 11: 63-66.
- Salter J.W. (1866) Appendix: On the fossils of North Wales. Memoir of the Geological Survey of Great Britain, 3: 240-381, pls 1-26.
- Schmidt W. & Geukens F. (1959) Nouveaux gîtes à Brachiopodes dans le Salmien inférieur du Massif de Stavelot. Bulletin de la Société belge de Géologie, de Paléontologie et d'Hydrologie, 67(2): 159-161.
- Servais T. & Sintubin M. (2013) Avalonia, Armorica, Perunica: terranes, microcontinents, microplates or palaeobiogeographical provinces? In: Bassett M.G. (Ed.) - Early Palaeozoic Peri-Gondwanan Terranes: New Insights from Tectonics and Biogeography Geological Society, London, Special Publications, 325: 103-115.
- Sheehan P.M. (1987) Late Ordovician (Ashgillian) brachiopods from the region of the Sambre and Meuse Rivers, Belgium. Bulletin de l'Institut royal des Sciences naturelles de

- Shellnutt J.G., Owen J.V., Yeh M.-W., Dostal J. & Nguyen D.T. (2019) - Long-lived association between Avalonia and the Meguma terrane deduced from zircon geochronology of metasedimentary granulites. *Scientific Reports*, 9(1): (4065). https://doi.org/10.1038/s41598-019-40673-9
- Sutton M.D., Bassett M.G. & Cherns L. (1999) Lingulate brachiopods from the Lower Ordovician of the Anglo-Welsh Basin. Part 1. *Monograph of the Palaeontographical Society*, London, 153(610): 1-60, pls 1–8.
- Sutton M.D., Bassett M.G. & Cherns L. (2000) Lingulate brachiopods from the Lower Ordovician of the Anglo-Welsh Basin. Part 2. *Monograph of the Palaeontographical Society*, London, 154(615): 61-114, pls 9–23.
- Vanguestaine M., Breuer P. & Lehnert O. (2004) Discovery of an Early Ordovician conodont fauna in the Salm Group of the Stavelot Inlier, Belgium. Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre, 74 (supplément): 39-48.
- Vanguestaine M. & Rushton A. (1979) Découverte d'un brachiopode inarticulé, Acrothele cf. bergeroni Walcott, dans le Revinien inférieur de Trois-Ponts, Cambrien du massif de Stavelot, Belgique. Annales de la Société géologique de Belgique, 102: 295-301.
- Verniers J., Herbosch A., Vanguestaine M., Geukens F, Delcambre B., Pingot J.-L., Belanger I., Hennebert M., Debacker T., Sintubin M. & De Vos W. (2002) - Cambrian-Ordovician-Silurian lithostratigraphic units (Belgium). *Geologica Belgica*, 4(1/2): 5-38.

- Waagen W. (1885) Salt Range fossils, vol. 1, part 4. Productus Limestone fossils, Brachiopoda. *Memoirs of the Geologi*cal Survey of India, Palaeontologia Indica, (series 13) fasc. 5: 729-770, pls 82–86.
- Walcott C.D. (1902) Cambrian Brachiopoda: Acrotreta, Linnarssonella; Obolus; with descriptions of new species. United States National Museum, Proceedings, 25: 577-612.
- Walcott C.D. (1908) Cambrian geology and paleontology, no. 3 – Cambrian Brachiopoda, descriptions of new genera and species: no. 4 – Classification and terminology of the Cambrian Brachiopoda. *Smithsonian Miscellaneous Collections*, 53: 53-165.
- Wang Wenhui & Servais T. (2015) A re-investigation of the *Rhabdinopora flabelliformis* fauna form the early Tremadocian '*Dictyonema*' Shale in Belgium. *Geologica Belgica*, 18(1): 66-77.
- Webby B.D., Cooper R.A., Bergström S.M. & Paris F. (2004) - Stratigraphic framework and time slices. In: Webby B.D., Paris F., Droser M. & Percival I. (Eds) - The Great Ordovician Biodiversification Event: 41-47. Columbia University Press, New York.
- Williams A., Carlson S.J., Brunton C.H.C., Holmer L.E. & Popov L.E. (1996) - A supra-ordinal classification of the Brachiopoda. *Philosophical Transactions of the Royal Society* of London (series B), 351: 1171-1193.
- Williams A. & Holmer L.E. (1992) Ornamentation and shell structure of acrotretoid brachiopods. *Palaeontology*, 35(3): 657-692.