


First annotated draft genomes of nonmarine ostracods (Ostracoda, Crustacea) with different reproductive modes

Patrick Tran Van,^{1,2,*} Yoann Anselmetti,^{3,†} Jens Bast,^{1,‡} Zoé Dumas,¹ Nicolas Galtier ,³ Kamil S. Jaron ,¹ Koen Martens,^{4,5} Darren J. Parker ,^{1,2} Marc Robinson-Rechavi ,^{1,2} Tanja Schwander,¹ Paul Simion ,^{3,6} and Isa Schön^{4,7,*}

¹Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

²Swiss Institute of Bioinformatics, Lausanne 1015, Switzerland

³ISEM—Institut des Sciences de l'Evolution, Montpellier 34090, France

⁴Royal Belgian Institute of Natural Sciences, OD Nature, Freshwater Biology, Brussels 1000, Belgium

⁵Department of Biology, University of Ghent, Ghent 9000, Belgium

⁶Université de Namur, LEGE, URBE, Namur 5000, Belgium

⁷University of Hasselt, Research Group Zoology, Diepenbeek 3590, Belgium

[†]Present address: CoBIUS Lab, Department of Computer Science, University of Sherbrooke, 2500 Boulevard de l'Université, Sherbrooke, QC J1K 2R1, Canada.

^{*}Present address: Institute for Zoology, University of Cologne, Köln, Germany.

*Corresponding authors: Department of Ecology and Evolution, University of Lausanne, Biophore, CH-1015 Lausanne, Switzerland. Patrick.tranvan@unil.ch (P.T.V.); RBINS, Vautierstraat 29, B-1000 Brussels, Belgium. ischoen@naturalsciences.be (I.S.)

Abstract

Ostracods are one of the oldest crustacean groups with an excellent fossil record and high importance for phylogenetic analyses but genome resources for this class are still lacking. We have successfully assembled and annotated the first reference genomes for three species of nonmarine ostracods; two with obligate sexual reproduction (*Cyprideis torosa* and *Notodromas monacha*) and the putative ancient asexual *Darwinula stevensoni*. This kind of genomic research has so far been impeded by the small size of most ostracods and the absence of genetic resources such as linkage maps or BAC libraries that were available for other crustaceans. For genome assembly, we used an Illumina-based sequencing technology, resulting in assemblies of similar sizes for the three species (335–382 Mb) and with scaffold numbers and their N50 (19–56 kb) in the same orders of magnitude. Gene annotations were guided by transcriptome data from each species. The three assemblies are relatively complete with BUSCO scores of 92–96. The number of predicted genes (13,771–17,776) is in the same range as Branchiopoda genomes but lower than in most malacostracan genomes. These three reference genomes from nonmarine ostracods provide the urgently needed basis to further develop ostracods as models for evolutionary and ecological research.

Keywords: ancient asexual; sexual; *Darwinula stevensoni*; *Cyprideis torosa*

Introduction

Relevance of ostracods

Ostracoda are small, bivalved crustaceans, widely occurring in almost all aquatic habitats as part of the meiobenthos and periphyton. There are 2330 formally described species of extant nonmarine ostracods (Meisch et al. 2019) and at least another 7000 described species of extant marine ostracod species (see Schön and Martens 2016 for an estimate by S. Brandao). Their calcified valves are preserved as microfossils, making them the extant arthropod group with the most extensive fossil record. The group has an estimated (Cambrian) age of c. 500 myr (millions of years) according to a molecular clock (Oakley et al. 2013), and c. 450 myr (Ordovician; Maddocks 1982) to 509 myr (Wolfe et al. 2016) according to the fossil record. This makes them one of the oldest extant pancrustacean groups (Figure 1). Because of their excellent fossil data, evolutionary events can be dated with real-time estimates making ostracods ideal models for evolutionary research (Butlin and

Menozzi 2000; Oakley and Cunningham 2002; Oakley et al. 2013; Schön and Martens 2016).

Contrary to the extensive focus on this group for palaeontological research, there is a total lack of published ostracod genomes, and even isolated genomic data from ostracods in open access databases are still rare. Thus, the only resources available beyond individual gene sequences are four mitogenomes [the marine ostracods *Vargula hilgendorffii* (Ogoh and Ohmiya 2004; GenBank accession number NC_005306) and *Cypridina dentata* (Wang et al. 2019; NC_042792); and two unpublished mitogenomes from *V. tsujii* (NC_039175) and *Cyprideis torosa* (PRJNA302529)]. Also, raw Illumina DNA sequencing reads of the podocopid ostracod *Eucypris virens* have been generated as part of a study testing DNA extraction methods for high-throughput sequencing in zooplankton (SRX8021019; Beninde et al. 2020) but these have neither been assembled nor annotated. In studies on crustacean phylogenies and gene expression (see Supplementary Table S1 for details), raw RNA-sequencing reads have been

Received: December 01, 2020. Accepted: February 5, 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of Genetics Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1 The phylogenetic position of the Ostracoda among the pancrustaceans and their age estimated from fossil and molecular data. Modified from [Oakley et al. \(2013\)](#). Different pancrustaceans are indicated by branches in different colors. The Ostracoda include the Podocopida, Platycopida, and Myodocopida. Here, three representatives of the Platycopida (indicated in purple) have been sequenced. The phylogenetic clade to which *D. stvensoni* belongs, is indicated by D, the clade to which *C. torosa* and *N. monacha* belong, is indicated by *. Black horizontal bars represent the range of age estimates in myr from Bayesian analyses by [Oakley et al. \(2013\)](#). The letters A–C in the black boxes indicated fossils that were used for calibrations of age estimates.

generated for a total of 12 species coming from the three major ostracod lineages (Mydocopida, Halocyprida, and Podocopida), but the number of assembled and annotated ostracod genes in these studies remains very limited, ranging between 4 and 822 genes.

Choice of model species

Extant nonmarine ostracods show a high prevalence of asexual reproduction ([Chaplin et al. 1994](#); [Butlin et al. 1998](#); [Martens et al. 1998](#)), which has evolved several times independently in different ostracod lineages and is most frequent in the Cyprididae and the Darwinulidae. Ostracods are thus an ideal group to further study the paradox of sex, which remains one of the most puzzling questions in evolutionary biology ([Bell 1982](#); [Otto and Lenormand 2002](#); [Schön et al. 2009a](#); [Neiman et al. 2018](#)). The most important

sets of hypotheses explaining why sex is advantageous despite its direct costs are based on the fact that physical linkage among loci generates different forms of selective interference (recently reviewed in [Otto 2020](#)). Genome-wide data are very valuable to test if asexuals indeed are affected by these predictions (e.g., [Glémin et al. 2019](#); [Jaron et al. 2020](#)) and to develop insights into mechanisms such as gene conversion ([Omilian et al. 2006](#)), DNA repair ([Schön and Martens, 1998](#); [Hecox-Lea and Mark Welch 2018](#)), or horizontal gene transfer ([Gladyshev et al. 2008](#); [Danchin et al. 2010](#); [Boschetti et al. 2012](#); [Paganini et al. 2012](#); [Flot et al. 2013](#)). Such data are also needed to further test for general consequences of asexuality beyond lineage-specific effects ([Jaron et al. 2020](#)). For many animal groups in which asexuality is frequent, genomic data are limited to a few representatives only ([Tvedte et al. 2019](#)) or are totally absent like in the Ostracoda.

Of all extant nonmarine ostracods, the Cyprididae (cyprids) are most speciose, comprising 42% of all known species (Meisch et al. 2019). They would thus be an obvious choice for genomic studies, also because in this ostracod family, mixed reproduction with sexual and asexual females and geographic parthenogenesis is very common (Horne et al. 1998). Asexual cyprids, however, are often polyploid (Adolfsson et al. 2010; Symonová et al. 2018), probably because of hybridization between males and asexual females through accidental mating (Schmit et al. 2013). Consequently, genome sizes are relatively large (Jeffery et al. 2017; Gregory 2020) up to 3.13 pg which equals more than 3 Gb. These features are likely to seriously complicate genomic assemblies and annotations in the absence of any genomic resources for ostracods, which is why we did not choose any asexual cyprid ostracods for this genome project. Instead, we have selected three other species of nonmarine ostracods, one putative ancient asexual darwinulid ostracod and two species with obligate sexual reproduction.

The ostracod family Darwinulidae is one of the two last remaining animal groups which are still supported as being genuine ancient asexuals (Heethoff et al. 2009; Schön et al. 2009b; Schwander 2016) and comprises about 35 morphospecies (Meisch et al. 2019). All darwinulids are brooders with valve dimorphisms between males and females that are detectable in the fossil record. Martens et al. (2003) showed that males have been absent in this family for at least 200 myr. One study reported a few males in a single darwinulid species (Smith et al. 2006) but proof of the functionality of these males for successful mating and meaningful genetic exchange could not be provided. Such (potential) atavistic males have also been reported in other putative ancient asexuals (Heethoff et al. 2009). The type species of the Darwinulidae, *Darwinula stevensoni*, has been asexual since c. 20 myr (Straub 1952), occurs on all continents except Antarctica (Schön et al. 2012) and in a wide range of habitats (Schön et al. 2009b). *Darwinula stevensoni* is the best investigated darwinulid ostracod so far and has been the subject of ecological (Van Doninck et al. 2002, 2003a, 2003b; Van den Broecke et al. 2013) and molecular research using DNA sequence data from single genes (Schön et al. 1998; 2003; Martens et al. 2005; Schön et al. 2012). These studies revealed that *D. stevensoni* is most likely apomictic or functionally mitotic (following the definition of apomixis in animals as in Schön et al. 2009a). The species also has low mutation rates as there appears to be no (Schön et al. 1998) or low (Schön and Martens 2003; Schön et al. 2009b) allelic divergence within individuals, and genetic differences between populations from different continents can be attributed to ancient vicariant processes (Schön et al. 2012). It has also been suggested that gene conversion is common in this species (Schön and Martens 1998; 2003), which could be an explanation for the low observed mutation rates. These results, however, were based on a limited number of genes and require further confirmation with genome-wide data. *Darwinula stevensoni* has a life cycle of 1 year in Belgium (Van Doninck et al. 2003b) and up to 4 years in more northern regions (McGregor 1969 in Northern America; Ranta 1979 in Finland), which is exceptionally long for a nonmarine ostracod. It can survive a wide range of temperatures, salinities (Van Doninck et al. 2002), and oxygen concentrations (Rossi et al. 2002). The total genome size of *D. stevensoni* has been estimated as 0.86–0.93 pg with flow cytometry (Pacziesniak, unpublished), approximating 900 Mb. There is no information on the ploidy level of *D. stevensoni*, except for the study by Tétart (1979) showing 22 dot-like chromosomes.

Because of its putative ancient asexuality, no close sexual relatives of *D. stevensoni* are available for comparative, genomic analyses. We have chosen two fully sexual nonmarine ostracod species from the Cytheroidea and the Notodromadidae with high population densities in Belgium as comparisons to the putative ancient asexual: *C. torosa* and *Notodromas monacha*, respectively. *Cyprideis torosa* inhabits brackish waters and is the only extant species of this genus in Europe (Meisch 2000). It has been the subject of various biological and especially palaeontological and geochemical studies (see for example, Heip, 1976a, 1976b; De Deckker et al. 1999; Keyser 2005). Frogley and Whittaker (2017) suggested that *C. torosa* is at least of Pleistocene origin (c. 2.5 myr) but might be older. There are only two molecular studies of this species based on single genes (Schön and Martens 2003; Schön et al. 2017). No information on the genome size or the karyotype of *C. torosa* is currently available.

The second sexual ostracod species analyzed here, *N. monacha*, occurs throughout the Northern hemisphere and is a nonmarine ostracod with a most peculiar behavior: it is partially hyponeustonic, hanging upside down attached to the water surface (Meisch 2000). The fossil record of *N. monacha* goes back to the Miocene (max 23 myr—Janz 1997), and its genome size is at 0.87 pg (Jeffery et al. 2017; Gregory 2020) very similar to that of *D. stevensoni*. This species has not yet been the subject of any molecular studies.

Our aim here is to provide the first reference genome data of nonmarine ostracods from three different species with varying reproductive modes: the putative ancient asexual *D. stevensoni* and the two obligate sexuals, *C. torosa* and *N. monacha*. We also generate transcriptomes of these species to facilitate genome annotations.

Materials and methods

Sample collection for genome and transcriptome sequencing

All three nonmarine ostracod species were sampled in Belgian lakes where previous research had shown that these species occurred (Schön and Martens 2003; Merckx et al. 2018). Living ostracods were sampled using a hand net with a mesh size of 150 μm . The hand net was swept in between the vegetation and forcefully right above the surface of the sediment for collecting *D. stevensoni* and *C. torosa*. *N. monacha* was sampled by moving the net on the water surface. Nonmarine ostracods were kept in habitat water. Their taxonomic identity was confirmed, and they were sorted alive under a binocular microscope as described by Martens and Horne (2016). Individual ostracods were picked with a pipette and transferred into sterilized EPA water in which they were maintained until DNA and RNA were extracted. More details on the origin of biological samples are provided in Supplementary Table S2.

For generating reference genomes, DNA was extracted from a single female of each species using the QIAamp DNA Micro kit according to the manufacturer's instructions. The extracted DNA from single females was amplified in two independent reactions using the SYNGIS TruePrime WGA kit and then pooled, to generate sufficient DNA for preparing different libraries. To generate transcriptomes for annotation of reference genomes, RNA was extracted from 40 pooled individuals per species from the same collection batch. For this, individuals were frozen in liquid nitrogen and, after addition of Trizol (Life Technologies), mechanically crushed with beads (Sigmund Lindner). Next, chloroform and ethanol-extraction methods were applied to the homogenized

tissue and the aqueous layer transferred to RNeasy MinElute Columns (Qiagen). Subsequent steps of RNA extraction were done following the RNeasy Mini Kit protocol, including DNase digestion. Finally, RNA was eluted into water and stored at -80°C . RNA quantity and quality were estimated with the NanoDrop (Thermo Scientific) and Bioanalyzer (Agilent).

Genome assembly

We prepared five genomic DNA libraries for each reference genome (three 2×125 bp paired-end libraries with average insert sizes of 250–300, 550 and 700 bp, and two mate-pair libraries with average insert sizes of 3000 and 5000 bp; see Supplementary Table S3 for more details) with the Illumina TruSeq DNA Library Prep Kit. Reads were generated with the Illumina HiSeq 3000 system for a total coverage between $351\times$ and $386\times$ (Supplementary Table S3).

Reads were filtered with Trimmomatic v0.36 (Bolger et al. 2014) and NxTrim v0.4.1 (O'Connell et al. 2015). Because of uneven coverage produced by PCR-based whole-genome amplification (Chen et al. 2013; Oyola et al. 2014), we first normalized reads using BMap v36.59 (Bushnell 2014) and then assembled into contigs with SPAdes v3.10.1 (Bankevich et al. 2012). Scaffolding was performed using SSPACE v3.0 (Boetzer et al. 2011). Scaffolds identified as contaminants were filtered out using Blobtools v1.0 (Laetsch and Blaxter 2017). The completeness of genomes assemblies was assessed with BUSCO v3.0.2 (Seppey et al. 2019) against the *arthropoda_odb9* dataset. More details of the assembly pipelines and the applied parameters can be found in Supplementary Material SM1.

Protein coding gene annotation

Libraries were prepared using the Illumina TruSeq Stranded RNA kit, following the manufacturer's instructions. RNA reads were generated with the Illumina HiSeq 2500 system (Supplementary Table S4). Reads were filtered with Trimmomatic v0.36. All trimmed reads were mapped against the genomes with STAR v2.5.3a (Dobin et al. 2013) and further assembled with Trinity v2.5.1 (Haas et al. 2013) under the "genome guided" mode to produce transcriptome assemblies.

The obtained transcriptomes and protein evidence were used to train and predict protein coding genes using MAKER v2.31.8 (Holt and Yandell 2011). Predicted protein coding genes were functionally annotated with Blast2GO v5.5.1 (Conesa et al. 2005; Götz et al. 2008) against the NCBI *non-redundant arthropods* protein database (v 2018-10).

More details of the annotation pipelines and the applied parameters can be found in Supplementary Material SM2.

GenomeScope analyses

The whole genome amplification approach, which we used in the present study because of the small body size of individual ostracods, generated unequal read coverage of ostracod genomes and prevented us from directly estimating genome sizes and levels of heterozygosity from the assemblies. To overcome this problem, we re-sequenced two individual ostracods each of *D. stevensoni* and *N. monacha* without whole genome amplification, preparing libraries with the NEBNext[®] Ultra[™] II DNA Library Prep Kit for Illumina. Reads were filtered with Trimmomatic v0.36 and analyzed using GenomeScope v2.0 (Ranallo-Benavidez et al. 2020) to correctly estimate genome size and heterozygosity. More details on the analyses are provided in the Supplementary Material SM3.

Table 1 Quality features of published crustacean genomic assemblies of the last 4 years and of the current study

Class	Order	Species	Size	No. of scaffolds	N50	BUSCO	Reference
Branchiopoda	Diplostraca	<i>Daphnia pulex</i> ^{a,b}	156	1,822	1,661	96	Ye et al. (2017)
Branchiopoda	Diplostraca	<i>D. magna</i> ^{a,b}	130	4,193	10,124	96.7 (C)	Lee et al. (2019)
Branchiopoda	Notostraca	<i>Lepidurus arcticus</i>	73	7,167	116	98.4 (C)	Savojardo et al. (2019)
Branchiopoda	Notostraca	<i>L. apus lubbocki</i>	90	20,738	402	97.8 (C)	Savojardo et al. (2019)
Branchiopoda	Spinicaudata	<i>Eulimnadia texanac</i>	120	112	18,000	n.i.	Baldwin-Brown et al. (2018)
Copepoda	Cyclopoida	<i>Apocyclops royi</i>	258	97,072	n.i.	50 (C)	Jørgensen et al. (2019)
Copepoda	Cyclopoida	<i>Oithona nana</i>	85	4,626	401	n.i.	Madoui et al. 2017
Copepoda	Harpacticoida	<i>Tigriopus californicus</i>	190	459	298	94.5 (C)	Barreto et al. (2018)
Copepoda	Harpacticoida	<i>T. japonicus</i>	197	339	10,650	96 (C)	Jeong et al. (2020)
Copepoda	Harpacticoida	<i>T. kingsejongensis</i>	295	270,823	159	61.1 (C)	Kang et al. 2017
Ostracoda	Podocopida	<i>Cyprideis torosa</i>	335	132,611	19	86.6 (C)	Current study
Ostracoda	Podocopida	<i>Darwinula stevensoni</i>	382	62,118	56	91.9 (C + F) 93.7 (C)	Current study
Ostracoda	Podocopida	<i>Notodromas monacha</i>	377	62,251	42	95.8 (C + F) 92.7 (C) 94.4 (C + F)	Current study
Malacostraca	Amphipoda	<i>Parhyale hawaiiensis</i> ^{b,d}	4,024	100,000	69	n.i.	Kao et al. (2016)
Malacostraca	Isopoda	<i>Armadillidium vulgare</i>	1,725	43,451	51	87.9 (C)	Chebbi et al. (2019)
Malacostraca	Decapoda	<i>Cherax quadricarinatus</i> ^c	3,237	508,682	33	81.3 (C)	Tan et al. (2020)
Malacostraca	Decapoda	<i>Eriocheir japonica sinensis</i> ^c	1,270	1,368	3,185	92.7 (C)	Tang et al. (2020)
Malacostraca	Decapoda	<i>Palaeomon carinicaudab</i>	9,185	28,089,718	586	n.i.	Li et al. (2019)
Malacostraca	Decapoda	<i>Penaeus monodon</i>	1,600	1,211,364	2	96.8 (C + F)	Van Quyen et al. (2020)
Malacostraca	Decapoda	<i>Litopenaeus vannameib</i> ^c	1,664	4,682	606	95	Zhang et al. (2019)
Malacostraca	Decapoda	<i>Marsupenaeus japonicus</i>	924	37,192,281	1	97	Yuan et al. (2018)
Malacostraca	Decapoda	<i>Procambarus virginialis</i>	3,300	3,752,011	39	n.i.	Gutkunst et al. (2018)

Assembly size is provided in million base pairs, scaffold N50 in kilo base pairs, and BUSCO scores in %. Letters behind BUSCO scores indicate the % of complete single copy genes (C) or % of single and fragmented single copy genes (C + F), respectively. Where BUSCO scores lack brackets, no further information on completeness of single copy genes was provided. n.i. = no information available.

^a Anchoring of scaffolds in existing genome assembly.

^b Linkage map available.

^c Long-read technology.

^d BAC library available.

Data availability

Raw sequence reads have been deposited in NCBI's sequence read archive under the following bioprojects: PRJNA515625 (reference genomes, Supplementary Table S3) and PRJNA631617 (RNA-seq for annotations and resequenced individuals, Supplementary Tables S4 and S5).

Genome assemblies and annotations have been deposited in the European Nucleotide Archive (ENA) under the accession number PRJEB38362 (Supplementary Tables S6 and S7). Codes for the analyses are available at: https://github.com/AsexGenomeEvol/Ostracoda_genomes. Additional supplementary material is available at the figshare collection page, including detailed descriptions of the methods (Supplementary Methods SM1–SM3), Tables S1 and S2, and S6 and S8, and the results of the GenomeScope analyses of *D. stevensoni* and *N. monacha* (Supplementary Figures S1, A–D). Supplemental Material available at figshare: <https://doi.org/10.25387/g3.13858817>.

Results and discussion

First ostracod reference genomes and their attributes

We produced the first *de novo* reference genomes of nonmarine ostracods, namely of the three species *D. stevensoni*, *C. torosa*, and *N. monacha* with different reproductive modes (see Supplementary Material SM1 and Tables S3 and S4 for more details on the assemblies). We used a whole genome amplification approach (WGA), because the TruSeq DNA Nano library prep kit for Illumina sequencing or low input protocols for PacBio (Duncan et al. 2019) were not available when these assemblies were generated. We would not recommend WGA for future studies because this PCR-based method generated uneven coverage, and consequently, problems for applying routine genome assembly methods and estimates of genome size and heterozygosity. Despite these limitations, our approach produced genome assemblies that are useful for future research as will be outlined below.

When assessing the quality of the obtained ostracod *de novo* genome assemblies, the assembly of the putative ancient asexual, *D. stevensoni*, had the best contiguity, with the largest N50 although the total number of scaffolds was similar to *N. monacha* (Table 1, Supplementary Table S6). The genome of the putative ancient asexual is furthermore the most complete as shown by its total BUSCO score of 96% and of 94% for complete single copy genes (Table 1). The quality of the genome from the obligate sexual ostracod *C. torosa* is the lowest of the three ostracod species as it has the highest number of scaffolds, and the lowest N50; it is also less complete with a total BUSCO score of 92% (Supplementary Table S7) and of 87% for complete single copy genes (Table 1). All three species have similar numbers of predicted genes and transcripts (Supplementary Table S7).

Ostracod genome sizes estimated with flow cytometry are somewhat larger than the estimates that we obtained here from GenomeScope analyses of re-sequenced individual ostracods. The haploid genome size of *D. stevensoni* was estimated at 420–455 Mb with flow cytometry (Pacziesniak, unpublished) while we estimated 362 Mb from sequence reads (Supplementary Figure S1, A and B). Similarly, the size of the haploid genome of *N. monacha* is estimated at 425 Mb with flow cytometry (Jeffery et al. 2017; Gregory 2020), which is larger than the 385 Mb (Supplementary Figure S1, C and D) that we obtained from sequence reads. It thus seems that either the genome size

estimates by flow cytometry are incorrect or that some parts of each genome are missing from our sequencing reads. Transposons and repeat-rich genomic regions can contribute to gaps in genomic assemblies (Peona et al. 2020). Some of these missing regions could also be GC rich, a feature which is known to cause a sequencing bias with Illumina technology (see for example, Chen et al. 2013, Botero-Castro et al. 2017). Acquiring more complete genome assemblies will require the additional application of long-read technologies to ostracods.

Genome-wide estimates of heterozygosity are especially interesting for asexual taxa because the absence of recombination is expected to cause accumulation of mutations, resulting in increasing allelic divergences within individuals (Birky 1996). Jaron et al. (2020) identified three factors driving intragenomic heterozygosity in asexuals: how the transition to parthenogenesis occurred, which cytological mechanism underlies parthenogenesis and how long asexual reproduction has been ongoing. Based on sequencing reads from individual ostracods, we estimate heterozygosity of the putative ancient asexual ostracod *D. stevensoni* to be 0.92%–0.99% (Supplementary Figure S1, A and B) and 1.32%–1.43% for the sexual *N. monacha* (Supplementary Figure S1, C and D). The genome-wide heterozygosity of *D. stevensoni* matches to some extent an earlier study on intra-individual divergence in three nuclear genes of *D. stevensoni* (Schön and Martens 2003). The finding of almost 1% heterozygosity in *D. stevensoni* is remarkable, given that all previous genome-wide estimates for asexual arthropods that did not evolve via hybridization revealed extremely low levels of heterozygosity (Jaron et al. 2020). Yet heterozygosity is clearly less than the estimates for parthenogenetic species with known hybrid origin (1.73%–8.5%) or polyploidy (1.84%–33.21%) (Jaron et al. 2020), supporting the view that *D. stevensoni* is neither a hybrid nor a polyploid. Asexual reproduction in ostracods is thought to be apomictic (Chaplin et al. 1994), implying that observed heterozygosity levels are largely dependent on the relative impact of heterozygosity losses from gene conversion and heterozygosity gains from new mutations. Given the apparent absence of sex and recombination for millions of years (Straub 1952), it is perhaps surprising that heterozygosity in this putative ancient asexual ostracod is not larger. This may suggest that genome-wide rates of gene conversion and mutation are comparable in this species.

Genome contiguity of ostracod assemblies as compared to other crustaceans

We here compare the qualities of our ostracod genome assemblies to those of 19 other crustacean species (Table 1) published in the last 4 years. We only include studies with complete assemblies and sufficient information to assess assembly qualities. We assessed the contiguity of the three *de novo* ostracod genome assemblies by the number of scaffolds and their N50. Both features are comparable to those of the copepod *Apocyclops royi* (Jørgensen et al. 2019) and the amphipod *Parhyale hawaiiensis* (Kao et al. 2016) (Table 1) and better than for crustaceans with larger genomes such as the decapods *Cherax quadricarinatus* (Tan et al. 2020), *Palaeomon carinicauda* (Li et al. 2019), *Penaeus mondon* (Van Quyen et al. 2020), *Marsupenaeus japonicus* (Yuan et al. 2018), and *Procamburus virginalis* (Gutekunst et al. 2018; Table 1). Genome assemblies of several other crustaceans, however, have smaller scaffold numbers and higher N50 and thus better contiguities than the assemblies obtained here for nonmarine ostracods. For the two notostracan *Lepidurus* species (Savojardo et al. 2019), this can probably be explained by their smaller genome sizes. For other crustaceans, genome assemblies or linkage maps have

been available beforehand which have considerably improved assembly qualities (Table 1) as in the examples of the cladoceran *Daphnia pulex* (Ye et al. 2017), *D. magna* (Lee et al. 2019), and the copepod *Tigriopus japonicus* (Jeong et al. 2020). No such genomic resources are currently available for ostracods. Finally, other studies of crustacean genomes with better assembly contiguities (the branchiopod *Eulimnadia texana*—Baldwin-Brown et al. 2018, and the decapod *Erichoir japonica sinensis*—Tang et al. 2020, the copepod *Tigriopus californicus*—Jeong et al. 2020, and the isopod *Armadillium vulgare*—Chebbi et al. 2019) have used a combination of Illumina and long-read technologies (Table 1). Long-read technologies such as PacBio used to require a relatively large amount of high-molecular weight DNA (Solares et al. 2018), which could not be obtained for ostracods with their very low yields of high-molecular weight DNA from individual specimens and their small body sizes as compared to many other crustaceans (Schön and Martens 2016). We hope that low input protocols for PacBio (Duncan et al. 2019) and other long-read technologies can be successfully applied to ostracods in the future, in which case the genome assemblies obtained here could form the basis for subsequent hybrid assemblies. Optimizing Oxford Nanopore Technology for nonmarine ostracods has already commenced (Schön et al. in prep.).

Genome annotations of ostracods and other crustaceans

Because our *de novo* ostracod genome assemblies are relatively complete (see BUSCO scores in Table 1), we will here also briefly compare some features of predicted protein coding genes with those of other crustaceans (Supplementary Table S8). We have predicted 13,771–17,776 protein coding genes in the three nonmarine ostracod genomes (Supplementary Tables S7 and S8), with the highest number for the sexual *C. torosa* and an intermediate estimate for the putative ancient asexual *D. stevensoni*. The number of annotated protein coding genes in nonmarine ostracods is similar to estimates for various branchiopods and the copepods *Oithona nana*, *Tigriopus californicus*, and *T. kingsejongensis* but lower than in most malacostracans (Supplementary Table S8). Not all genome studies of crustaceans cited here contain information on other features of coding genes, such as the average size of genes, introns, and exons (Supplementary Table S7). Comparisons of these features are therefore limited and will not be further discussed here but we provide available data of these features for ostracods and other crustacean genomes for reference.

Gene annotation in general but especially in the crustaceans is challenging; this is for example illustrated by the much lower numbers of protein coding genes (18,440) which are predicted in the novel reference genome of the cladoceran *Daphnia pulex* by Ye et al. (2017) as compared to the first assembly of *D. pulex* with more than 30,000 predicted genes (Colbourne et al. 2011). Even more difficult is assigning gene functions to annotated crustacean genomes (Rotlantz et al. 2018). The novel data on predicted genes and transcripts from nonmarine ostracods in the current study will significantly contribute to future genome annotations in crustaceans and other arthropods. The genes and transcripts predicted here can also provide the baseline for future gene expression studies of nonmarine and marine ostracods.

Conclusions

We have successfully obtained *de novo* genome assemblies for three species of nonmarine ostracods with different reproductive

modes. These represent the first quality reference genomes for ostracods. Given the paucity of genome assemblies from crustaceans as compared to insects or other arthropods, these assemblies are important tools to further develop ostracods as models for evolutionary and ecological research, also including marine species. Even if the *de novo* genome assemblies are somewhat fragmented and not yet at the chromosome level, they have a high level of completeness and will thus facilitate future studies of ostracods. The genomes presented here can also provide the first step toward a genomic assessment of the putative ancient asexual status of nonmarine darwinulid ostracod species.

Acknowledgments

Marie Cours, Tijs Van Den Bergen, and Jeroen Venderickx are acknowledged for technical support in sampling and sorting ostracod samples. We also thank Kristiaan Hoedemakers and Jeroen Venderickx for their assistance in finalizing the figure.

Funding

This research was funded by Belgian Federal Science Policy (BR/314/PI/LATTECO) and a grant from the Swiss National Science Foundation (CRSII3_160723).

Conflicts of interest: None declared.

Literature cited

- Adolfsson S, Michalakis Y, Paczesniak D, Bode SNS, Butlin RK, et al. 2010. Evaluation of elevated ploidy and asexual reproduction as alternative explanations for geographic parthenogenesis in *Eucypris virens* ostracods. *Evolution* 64:986–997. doi: 10.1111/j.1558-5646.2009.00872.x.
- Baldwin-Brown JG, Weeks SC, Long AD. 2018. A new standard for crustacean genomes: the highly contiguous, annotated genome assembly of the clam shrimp *Eulimnadia texana* reveals HOX gene order and identifies the sex chromosome. *Genome Biol Evol.* 10: 143–156. doi: 10.1093/gbe/evx280.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, et al. 2012. Spades: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 19:455–477. doi: 10.1089/cmb.2012.0021.
- Barreto FS, Watson ET, Lima TG, Willett CS, Edmands S, et al. 2018. Genomic signatures of mitonuclear coevolution across populations of *Tigriopus californicus*. *Nat Ecol Evol.* 2:1250–1257. doi: 10.1038/s41559-018-0588-1.
- Bell G. 1982. *The Masterpiece of Nature: The Evolution and Genetics of Sexuality*. Berkeley, CA: University of California Press.
- Beninde J, Möst M, Meyer A. 2020. Optimized and affordable high-throughput sequencing workflow for preserved and nonpreserved small zooplankton specimens. *Mol Ecol Res.* 20: 1632–1646. doi: 10.1111/1755-0998.13228.
- Birky CW, Jr. 1996. Heterozygosity, heteromorphy, and phylogenetic trees in asexual eukaryotes. *Genetics* 144:427–437.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27:578–579. doi: 10.1093/bioinformatics/btq683.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. doi: 10.1093/bioinformatics/btu170.

- Boschetti C, Carr A, Crisp A, Eyres I, Wang-Koh Y, et al. 2012. Biochemical diversification through foreign gene expression in bdelloid rotifers. *PLoS Genet.* 8:e1003035. doi: 10.1371/journal.pgen.1003035.
- Botero-Castro F, Figuet E, Tilak MK, Nabholz B, Galtier N. 2017. Avian genomes revisited: Hidden genes uncovered and the rates versus traits paradox in birds. *Mol Biol Evol.* 34:3123–3131. doi: 10.1093/molbev/msx236.
- Bushnell B. 2014. BMAP: A fast, accurate, splice-aware aligner. 9th Annual Genomics of Energy & Environment Meeting, Walnut Creek, CA, USA.
- Butlin RK, Menozzi P. 2000. Open questions in evolutionary ecology: do ostracods have the answers? In: Horne DJ, Martens K (eds.), *Evolutionary Biology and Ecology of Ostracoda*. Dordrecht: Springer. p. 1–14. doi: 10.1023/A:1003925705735.
- Butlin RK, Schön I, Martens K. 1998. Asexual reproduction in non-marine ostracods. *Heredity* 81:473–480. doi: 10.1046/j.1365-2540.1998.00454.x.
- Chaplin JA, Havel JE, Hebert PDN. 1994. Sex and ostracods. *Trends Ecol Evol.* 9:435–439. doi: 10.1016/0169-5347(94)90127-9.
- Chebbi MA, Becking T, Moumen B, Giraud I, Gilbert C, et al. 2019. The genome of *Armadillidium vulgare* (Crustacea, Isopoda) provides insights into sex chromosome evolution in the context of cytoplasmic sex determination. *Mol Biol Evol.* 36:727–741. doi: 10.1093/molbev/msz010.
- Chen Y-C, Liu T, Yu C-H, Chiang T-Y, Hwang C-C. 2013. Effects of gc bias in next-generation-sequencing data on *de novo* genome assembly. *PLoS One* 8:e62856. doi: 10.1371/journal.pone.0062856.
- Colbourne JK, Pfrender ME, Gilbert D, Thomas WK, Tucker A, et al. 2011. The ecoresponsive genome of *Daphnia pulex*. *Science* 331: 555–561. doi: 10.1126/science.1197761.
- Conesa A, Götz S, García-Gómez JM, Terol J, Talon M, et al. 2005. Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21: 3674–3676. doi: 10.1093/bioinformatics/bti610.
- Danchin EG, Rosso MN, Vieira P, de Almeida-Engler J, Coutinho PM, et al. 2010. Multiple lateral gene transfers and duplications have promoted plant parasitism ability in nematodes. *Proc Natl Acad Sci USA.* 107:17651–17656. doi: 10.1073/pnas.1008486107.
- De Deckker P, Chivas AR, Shelley JMG. 1999. Uptake of Mg and Sr in the euryhaline ostracod *Cyprideis* determined from in vitro experiments. *Palaeogeogr Palaeoclimatol Palaeoecol.* 148: 105–116. doi: 10.1016/S0031-0182(98)00178-3.
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, et al. 2013. STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15–21. doi: 10.1093/bioinformatics/bts635.
- Duncan T, Kingan SB, Lambert CC, Baybayan P, Korf J. 2019. A low DNA input protocol for high-quality PacBio *de novo* genome assemblies. *J Biomol Technol.* 30:S1–S2.
- Flot J-F, Hespeels B, Li X, Noel B, Arkhipova I, et al. 2013. Genomic evidence for ameiotic evolution in the bdelloid rotifer *Adineta vaga*. *Nature* 500:453–457. doi: 10.1038/nature12326.
- Frogley MR, Whittaker JE. 2017. On the origins of *Cyprideis torosa* (Jones, 1850) and a short biography of Professor T. R. Jones. *J Micropalaeontol.* 36:8–12. doi: 10.1144/jmpaleo2015-023.
- Gladyshev EA, Meselson M, Arkhipova IR. 2008. Massive horizontal gene transfer in bdelloid rotifers. *Science* 320:1210–1213. doi: 10.1126/science.1156407.
- Glémin S, François CM, Galtier N. 2019. Genome evolution in outcrossing vs. selfing vs. asexual species. In: Anisimova M (ed.), *Evolutionary Genomics*. New York, NY: Humana. p. 331–369. doi: 10.1007/978-1-4939-9074-0_11.
- Götz S, Garcia-Gomez JM, Terol J, Williams TD, Nagaraj SH, et al. 2008. High-throughput functional annotation and data mining with the Blast2GO suite. *Nucleic Acids Res.* 36:3420–3435. doi: 10.1093/nar/gkn176.
- Gregory TR. 2020. Animal genome size database. Available online: <http://genomesize.com> (Accessed: 2020 October 7).
- Gutekunst J, Andriantsoa R, Falckenhayn C, Hanna K, Stein W, et al. 2018. Clonal genome evolution and rapid invasive spread of the marbled crayfish. *Nat Ecol Evol.* 2:567–573. doi: 10.1038/s41559-018-0467-9.
- Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, et al. 2013. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat Protoc.* 8:1494–1512. doi: 10.1038/nprot.2013.084.
- Hecox-Lea BJ, Mark Welch DB. 2018. Evolutionary diversity and novelty of DNA repair genes in asexual Bdelloid rotifers. *BMC Evol Biol.* 18:177. doi: 10.1186/s12862-018-1288-9.
- Heethoff M, Norton RA, Scheu S, Maraun M. 2009. Parthenogenesis in oribatid mites (Acari, Oribatida): evolution without sex. In: I Schön, K Martens, P Van Dijk, editor. *Lost Sex. The Evolutionary Biology of Parthenogenesis*. Dordrecht, The Netherlands: Springer Academic Publishers. p. 241–258. doi: 10.1007/978-90-481-2770-2_12.
- Heip C. 1976a. The life-cycle of *Cyprideis torosa* (Crustacea, Ostracoda). *Oecologia* 24:229–245. doi: 10.1007/BF00345475.
- Heip C. 1976b. The spatial pattern of *Cyprideis torosa* (Jones, 1850) (Crustacea: Ostracoda). *J Mar Biol Assoc.* 56:179–189. doi: 10.1017/S002531540002052X.
- Holt C, Yandell M. 2011. MAKER2: An annotation pipeline and genome-database management tool for second-generation genome projects. *BMC Bioinformatics.* 12:491. doi: 10.1186/1471-2105-12-491.
- Horne DJ, Baltanas A, Paris, G. 1998. Geographical distribution of reproductive modes in living non-marine ostracods. In: K Martens, editor. *Sex and Parthenogenesis. Evolutionary Ecology of Reproductive Modes in Non-Marine Ostracods*. Leiden: Backhuys Publishers. p. 77–99.
- Janz H. 1997. Die Ostrakoden der kleini-Schichten des miozänen Kratersees von Steinheim am Albuch (Süddeutschland). *Stutt Beitr Naturkd B* 251:1–101.
- Jaron KS, Bast J, Novell RW, Ranallo-Benavidez TR, Robinson-Rechavi M, et al. 2020. Genomic features of asexual animals. *J Hered.* esaa031. doi: 10.1093/jhered/esaa031.
- Jeffery NW, Ellis EA, Oakley TH, Gregory TR. 2017. The genome sizes of ostracod crustaceans correlate with body size and evolutionary history, but not environment. *J Hered.* 108:701–706. doi: 10.1093/jhered/esx055.
- Jeong CB, Lee BY, Choi BS, Kim MS, Park JC, et al. 2020. The genome of the harpacticoid copepod *Tigriopus japonicus*: Potential for its use in marine molecular ecotoxicology. *Aquat Toxicol.* 222:105462. doi: 10.1016/j.aquatox.2020.105462.
- Jørgensen TS, Nielsen BLH, Petersen B, Browne PD, Hansen BW, et al. 2019. The whole genome sequence and mRNA transcriptome of the tropical cyclopoid copepod *Apocyclops royi*. *G3 (Bethesda)* 9: 1295–1302. doi: 10.1534/g3.119.400085.
- Kang S, Ahn D-H, Lee J H, Lee S G, Shin S C, et al. 2017. The genome of the Antarctic-endemic copepod, *Tigriopus kingsejongensis*. *Gigascience.* 6:1–9.
- Kao D, Lai AG, Stamatakis E, Rosic S, Konstantinides N, et al. 2016. The genome of the crustacean *Parhyale hawaiiensis*, a model for animal development, regeneration, immunity and lignocellulose digestion. *elife* 5:e20062. doi: 10.7554/eLife.20062.

- Keyser D. 2005. Histological peculiarities of the nodding process in *Cyprideis torosa* (Jones) (Crustacea, Ostracoda). *Hydrobiologia* 538: 95–106. doi: 10.1007/s10750-004-4940-x.
- Laetsch DR, Blaxter ML. 2017. BlobTools: Interrogation of genome assemblies. *F1000Research* 6:1287. doi: 10.12688/f1000research.12232.1.
- Lee BY, Choi BS, Kim MS, Park JC, Jeong CB, et al. 2019. The genome of the freshwater water flea *Daphnia magna*: A potential use for freshwater molecular ecotoxicology. *Aquat Toxicol.* 210:69–84. doi: 10.1016/j.aquatox.2019.02.009.
- Li J, Lv J, Liu P, Chen P, Wang J, et al. 2019. Genome survey and high-resolution backcross genetic linkage map construction of the ridgetail white prawn *Exopalaemon carinicauda* applications to QTL mapping of growth traits. *BMC Genomics* 20:598. doi: 10.1186/s12864-019-5981-x.
- Maddocks R. 1982. Part 4. Ostracoda. In LG Abele, editor. *The Biology of Crustacea*. Vol 1: Systematics, the Fossil Record and Biogeography. Academic Press. p. 221–239.
- Madoui M-A, Poulain J, Sugier K, Wessner M, Noel B, et al. 2017. New insights into global biogeography, population structure and natural selection from the genome of the epipelagic copepod *Oithona*. *Mol Ecol.* 26:4467–4482.
- Martens K, Horne DJ. 2016. Collecting and processing living non-marine ostracods. *J Crust Biol.* 36:849–854. doi: 10.1163/1937240X-00002488.
- Martens K, Horne DJ, Griffiths HI. 1998. Age and diversity of non-marine ostracods. In: K Martens, editor. *Sex and Parthenogenesis. Evolutionary Ecology of Reproductive Modes in Non-Marine Ostracods*. Leiden: Backhuys Publishers. p. 37–55.
- Martens K, Rossetti G, Butlin RK, Schön I. 2005. Molecular and morphological phylogeny of the ancient asexual Darwinulidae (Crustacea, Ostracoda). *Hydrobiologia* 538:153–165. doi: 10.1007/PL00021867.
- Martens K, Rossetti G, Horne D. 2003. How ancient are ancient asexuals? *Proc R Soc Lond B* 270:723–729. doi: 10.1098/rspb.2002.2270.
- McGregor DL. 1969. The reproductive potential, life history and parasitism of the freshwater Ostracods *Darwinula stevensoni* (Brady and Robertson). In: JW Neale, editor. *The Taxonomy, Morphology and Ecology of Recent Ostracoda*. Edinburgh: Oliver and Boyd. p. 194–221.
- Meisch C. 2000. *Freshwater Ostracoda of Western and Central Europe*. Heidelberg, Germany: Springer.
- Meisch C, Smith RJ, Martens K. 2019. A subjective global checklist of the extant non-marine Ostracoda (Crustacea). *EJT* 25. doi: 10.5852/ejt.2019.492.
- Merckx T, Souffreau C, Kaiser A, Baardsen LF, Backeljau T, et al. 2018. Body size shifts in aquatic and terrestrial urban communities. *Nature* 558:113–116. doi: 10.1038/s41586-018-0140-0.
- Neiman M, Meirmans PG, Schwander T, Meirmans S. 2018. Sex in the wild: how and why field-based studies contribute to solving the problem of sex. *Evolution* 72:1194–1203. doi: 10.1111/evo.13485.
- O'Connell J, Schulz-Trieglaff O, Carlson E, Hims MM, Gormley NA, et al. 2015. NxTrim: optimized trimming of Illumina mate pair reads. *Bioinformatics* 31:2035–2037. doi: 10.1093/bioinformatics/btv057.
- Oakley TH, Cunningham CW. 2002. Molecular phylogenetic evidence for the independent evolutionary origin of an arthropod compound eye. *Proc. Natl Acad Sci USA.* 99:1426–1430. doi: 10.1073/pnas.032483599.
- Oakley TH, Wolfe JM, Lindgren AR, Zaharoff AK. 2013. Phylotranscriptomics to bring the understudied into the fold: monophyletic Ostracoda, fossil placement, and pancrustacean phylogeny. *Mol Biol E* 30:215–233. doi: 10.1093/molbev/mss216.
- Ogoh K, Ohmiya Y. 2004. Complete mitochondrial DNA sequence of the sea-firefly, *Vargula hilgendorffii* (Crustacea, Ostracoda) with duplicate control regions. *Gene* 327:131–139. doi: 10.1016/j.gene.2003.11.011.
- Omilian AR, Cristescu ME, Dudycha JL, Lynch M. 2006. Asexual recombination in asexual lineages of *Daphnia*. *Proc Natl Acad Sci USA.* 103:18638–18643. doi: 10.1073/pnas.0606435103.
- Otto SP. 2020. Selective interference and the evolution of sex. *J Hered.* esaa026. doi: 10.1093/jhered/esaa026.
- Otto SP, Lenormand T. 2002. Resolving the paradox of sex and recombination. *Nat Rev Genet.* 3:252–261. doi: 10.1038/nrg761.
- Oyola SO, Manske M, Campino S, Claessens A, Hamilton WL, et al. 2014. Optimized whole-genome amplification strategy for extremely AT-biased template. *DNA Res.* 21:661–671. doi: 10.1093/dnares/dsu028.
- Paganini J, Campan-Fournier A, Da Rocha M, Gouret P, Pontarotti P, et al. 2012. Contribution of lateral gene transfers to the genome composition and parasitic ability of root-knot nematodes. *PLoS One* 7:e50875. doi: 10.1371/journal.pone.0050875.
- Peona V, Blom MP, Xu L, Burri R, Sullivan S, et al. 2020. Identifying the causes and consequences of assembly gaps using a multiplatform genome assembly of a bird-of-paradise. *Mol Ecol Res.* doi: 10.1111/1755-0998.13252.
- Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. *Nat Commun.* 11:1–10. doi: 10.1038/s41467-020-14998-3.
- Ranta E. 1979. Population biology of *Darwinula stevensoni* (Crustacea, Ostracoda) in an oligotrophic lake. *Ann Zool Fenn.* 16:28–35. <https://www.jstor.org/stable/2373370>.
- Rossi V, Todeschi EBA, Gandolfi A, Invidia M, Menozzi P. 2002. Hypoxia and starvation tolerance in individuals from a riverine and a lacustrine population of *Darwinula stevensoni* (Crustacea: Ostracoda). *Arch Hydrobiol.* 154:151–171. doi: 10.1127/archiv-hydrobiol/154/2002/151.
- Rotllant G, Palero F, Mather PB, Bracken-Grissom HD, Santos MB. 2018. Preface: recent advances in crustacean genomics. *Hydrobiologia* 825:1–4. doi: 10.1007/s10750-018-3773-y.
- Savojardo C, Luchetti A, Martelli PL, Casadio R, Mantovani B. 2019. Draft genomes and genomic divergence of two *Lepidurus* tadpole shrimp species (Crustacea, Branchiopoda, Notostraca). *Mol Ecol Resour.* 19:235–244. doi: 10.1111/1755-0998.12952.
- Schmit O, Fukova L, Vandekerckhove J, Michalakakis Y, Matzke-Karasz R, et al. 2013. Mate recognition as a reproductive barrier in sexual and parthenogenetic *Eucypris virens* (Crustacea, Ostracoda). *Anim Behav.* 85:977–985. doi: 10.1016/j.anbehav.2013.02.021.
- Schön I, Butlin RK, Griffiths HI, Martens K. 1998. Slow molecular evolution in an ancient asexual ostracod. *Proc R Soc Lond B* 265: 235–242. doi: 10.1098/rspb.1998.0287.
- Schön I, Halse S, Martens K. 2017. *Cyprideis* (Crustacea, Ostracoda) in Australia. *J Micropalaeontol.* 36:31–37. doi: 10.1144/jmpaleo2016-032.
- Schön I, Martens K. 1998. Opinion: DNA repair in an ancient asexual - a new solution for an old problem? *J Nat Hist.* 32:943–948. doi: 10.1080/00222939800770481.
- Schön I, Martens K. 2003. No slave to sex. *Proc R Soc Lond B* 270: 827–833. doi: 10.1098/rspb.2002.2314.
- Schön I, Martens K. 2016. Ostracod (Ostracoda, Crustacea) genomics - Promises and challenges. *Mar Genomics* 29:19–25. doi: 10.1016/j.margen.2016.03.008.
- Schön I, Martens K, Van Dijk P, editors. 2009a. *Lost Sex. The Evolutionary Biology of Parthenogenesis*. Dordrecht, The Netherlands: Springer Academic Publishers.

- Schön I, Martens K, Van Doninck K, Butlin RK. 2003. Evolution in the slow lane: molecular rates of evolution in sexual and asexual ostracods (Crustacea: Ostracoda). *Biol J Linn Soc.* 79:93–100. doi: 10.1046/j.1095-8312.2003.00186.x.
- Schön I, Pinto RL, Halse S, Smith AJ, Martens K, et al. 2012. Cryptic species in putative ancient asexual Darwinulids (Crustacea, Ostracoda). *PLoS One.* 7:e39844. doi: 10.1371/journal.pone.0039844.
- Schön I, Rossetti G, Martens K. 2009b. Ancient asexual darwinulids: ancient scandals or scandalous gossip? In: I Schön, K Martens, P Van Dijk, editors. *Lost Sex. The Evolutionary Biology of Parthenogenesis.* Dordrecht, The Netherlands: Springer Academic Publishers, p. 217–240. doi: 10.1007/978-90-481-2770-2_11.
- Schwander T. 2016. The end of an ancient asexual scandal. *Curr Biol.* 26:R233–R235. doi: 10.1016/j.cub.2016.01.034.
- Seppy M, Manni M, Zdobnov EM. 2019. Busco: Assessing genome assembly and annotation completeness. *Methods Mol Biol.* 1962: 227–245. doi: 10.1007/978-1-4939-9173-0_14.
- Smith R, Kamiya T, Horne DJ. 2006. Living males of the 'ancient asexual' Darwinulidae (Ostracoda: Crustacea). *Proc R Soc B.* 273: 1569–1578. doi: 10.1098/rspb.2005.3452.
- Solares EA, Chakraborty M, Miller DE, Kalsow S, et al. 2018. Rapid low-cost assembly of the *Drosophila melanogaster* reference genome using low-coverage, long-read sequencing. *G3 (Bethesda)* 8:3143–3154. doi: 10.1534/g3.118.200162.
- Straub EB. 1952. Mikropaläontologische Untersuchungen im Tertiär zwischen Ehingen und Ulm a.d. Donau *Geol Jahrb.* 66:433–523.
- Symonová R, Vrbová I, Lamatsch DK, Paar J, Matzke-Karasz R, et al. 2018. Karyotype variability and inter-population genomic differences in freshwater ostracods (Crustacea) showing geographical parthenogenesis. *Genes* 9:150. doi: 10.3390/genes9030150.
- Tan MH, Gan HM, Lee YP, Grandjean F, Croft LJ, et al. 2020. A giant genome for a giant crayfish (*Cherax quadricarinatus*) with insights into *cox1* pseudogenes in decapod genomes. *Front Genet.* 11:201. doi: 10.3389/fgene.2020.00201.
- Tang B, Wang Z, Liu Q, Zhang H, Jiang S, et al. 2020. High-quality genome assembly of *Eriocheir japonica sinensis* reveals its unique genome evolution. *Front Genet.* 10:1340. doi: 10.3389/fgene.2019.01340.
- Tétart J. 1979. Les garnitures chromosomiques des ostracodes d'eau douce. *Trav Lab Hydrobiol.* 69–70:113–140. (In French).
- Tvedte ES, Logsdon JM, Jr., Forbes AA. 2019. Sex loss in insects: causes of asexuality and consequences for genomes. *Curr Opin Insect Sci.* 31:77–83. doi: 10.1016/j.cois.2018.11.007.
- Van den Broecke L, Vanfleteren J, Martens K, Schön I. 2013. Hurdles in investigating UVB damage in the putative ancient asexual *Darwinula stevensoni* (Ostracoda, Crustacea). *Br J Zool.* 143: 106–118.
- Van Doninck K, Schön I, De Bruyn L, Martens K. 2002. A general purpose genotype in an ancient asexual. *Oecologia* 132:205–212. doi: 10.1007/s00442-002-0939-z.
- Van Doninck K, Schön I, Maes F, De Bruyn L, Martens K. 2003a. Ecological strategies in the ancient asexual animal group Darwinulidae. *Freshwater Biol.* 48:1285–1294. doi: 10.1046/j.1365-2427.2003.01078.x.
- Van Doninck K, Schön I, Martens K, Goddeeris B. 2003b. The life cycle of the ancient asexual ostracod *Darwinula stevensoni* (Brady & Robertson, 1870) (Crustacea, Ostracoda) in a temperate pond. *Hydrobiologia* 500:331–340. doi: 10.1023/A:1024656920904.
- Van Quyen D, Gan HM, Lee YP, Nguyen DD, Nguyen TH, et al. 2020. Improved genomic resources for the black tiger prawn (*Penaeus monodon*). *Marine Genom.* 52:100751. doi: 10.1016/j.margen.2020.100751.
- Wang X, Xu Q, Xiao J, Miao X, Liu P, et al. 2019. First record of the complete mitochondrial genome of *Cypridina dentata* (Myodocopida: Cypridinidae). *Mitochondrial DNA B Resourc.* 4:1607–1608. doi: 10.1080/23802359.2019.1604097.
- Wolfe JM, Daley AC, Legg DA, Edgecombe GD. 2016. Fossil calibrations for the arthropod Tree of Life. *Earth Sci Rev.* 160:43–110. doi: 10.1016/j.earscirev.2016.06.008.
- Ye Z, Xu S, Spitze K, Asselman J, Jiang X, et al. 2017. A new reference genome assembly for the microcrustacean *Daphnia pulex*. *G3 (Bethesda)* 7:1405–1416. doi: 10.1534/g3.116.038638.
- Yuan J, Zhang X, Liu C, Yu Y, Wei J, et al. 2018. Genomic resources and comparative analyses of two economical penaeid shrimp species, *Marsupenaeus japonicus* and *Penaeus monodon*. *Mar Genomics* 39:22–25. doi: 10.1016/j.margen.2017.12.006.
- Zhang X, Yuan J, Sun Y, Li S, Gao Y, et al. 2019. Penaeid shrimp genome provides insights into benthic adaptation and frequent molting. *Nat Commun.* 10:1–14. Doi: 10.1038/s41467-018-08197-4.

Communicating editor: B. J. Andrews