

EDITORIAL

The taxonomic impediment: a shortage of taxonomists, not the lack of technical approaches

MICHAEL S. ENGEL^{1,‡,◉}, LUIS M. P. CERÍACO^{2,‡}, GIMO M. DANIEL^{3,‡,◉}, PABLO M. DELLAPÉ^{4,‡}, IVAN LÖBL^{5,‡}, MILEN MARINOV^{6,‡,*}, ROBERTO E. REIS^{7,‡}, MARK T. YOUNG^{8,‡}, ALAIN DUBOIS^{9,‡}, ISHAN AGARWAL¹⁰, PABLO LEHMANN A.¹¹, MABEL ALVARADO¹², NADIR ALVAREZ⁵, FRANCO ANDREONE¹³, KATYUSCIA ARAUJO-VIEIRA¹⁴, JOHN S. ASCHER¹⁵, DÉLIO BAÊTA¹⁶, DIEGO BALDO¹⁷, SUZANA A. BANDEIRA¹⁸, PHILLIP BARDEN¹⁹, DIEGO A. BARRASSO²⁰, LEILA BENDIFALLAH²¹, FLÁVIO A. BOCKMANN²², WOLFGANG BÖHME²³, ART BORKENT²⁴, CARLOS R. F. BRANDÃO²⁵, STEPHEN D. BUSACK²⁶, SETH M. BYBEE²⁷, ALAN CHANNING²⁸, STYLIANOS CHATZIMANOLIS²⁹, MAARTEN J. M. CHRISTENHUSZ³⁰, JORGE V. CRISCI³¹, GUILLERMO D'ELÍA³², LUIS M. DA COSTA³³, STEVEN R. DAVIS³⁴, CARLOS ALBERTO S. DE LUCENA³⁵, THIERRY DEUVE⁹, SARA FERNANDES ELIZALDE³⁶, JULIÁN FAIVOVICH³⁷, HARITH FAROOQ³⁸, ADAM W. FERGUSON³⁹, SPARTACO GIPPOLITI⁴⁰, FRANCISCO M. P. GONÇALVES⁴¹, VICTOR H. GONZALEZ⁴², ELI GREENBAUM⁴³, ISMAEL A. HINOJOSA-DÍAZ⁴⁴, IVAN INEICH⁹, JIANPING JIANG⁴⁵, SIH KAHONO⁴⁶, ADRIANO B. KURY⁴⁷, PAULO H. F. LUCINDA⁴⁸, JOHN D. LYNCH⁴⁹, VALÉRY MALÉCOT⁵⁰, MARIANA P. MARQUES⁵¹, JOHN W. M. MARRIS⁵², RYAN C. MCKELLAR⁵³, LUIS F. MENDES⁵¹, SILVIO S. NIHEI⁵⁴, KANTO NISHIKAWA⁵⁵, ANNEMARIE OHLER⁹, VICTOR G. D. ORRICO⁵⁶, HIDETOSHI OTA⁵⁷, JORGE PAIVA⁵⁸, DIOGO PARRINHA⁵¹, OLIVIER S. G. PAUWELS⁵⁹, MARTÍN O. PEREYRA⁶⁰, LUEJI B. PESTANA⁶¹, PAULO D. P. PINHEIRO⁵⁴, LORENZO PRENDINI³⁴, JAKUB PROKOP⁶², CLAUS RASMUSSEN⁶³, MARK-OLIVER RÖDEL⁶⁴, MIGUEL TREFAUT RODRIGUES⁵⁴, SARA M. RODRÍGUEZ⁶⁵, HEARTY SALATNAYA⁶⁶, ÍRIS SAMPAIO², ALBA SÁNCHEZ-GARCÍA⁶⁷, MOHAMED A. SHEBL⁶⁸, BRUNA S. SANTOS⁶⁹, MÓNICA M. SOLÓRZANO-KRAEMER⁷⁰, ANA C. A. SOUSA⁷¹, PAVEL STOEVT⁷², PABLO TETA⁷³, JEAN-FRANÇOIS TRAPE⁷⁴, CARMEN VAN-DÚNEM DOS SANTOS⁷⁵, KARTHIKEYAN VASUDEVAN⁷⁶, COR J. VINK⁷⁷, GERNOT VOGEL⁷⁸, PHILIPP WAGNER⁷⁹, TORSTEN WAPPLER⁸⁰, JESSICA L. WARE⁸¹, SONJA WEDMANN⁸² and CHIFUNDERA KUSAMBA ZACHARIE⁸³

¹*Division of Entomology, Natural History Museum, University of Kansas, Lawrence, Kansas, USA*

²*Museu de História Natural e da Ciência da Universidade do Porto, Porto, Portugal*

³*Department of Terrestrial Invertebrates, The National Museum, Bloemfontein, South Africa; Department of Biological & Environmental Sciences, Walter Sisulu University, Mthatha, South Africa*

⁴*División Entomología, Universidad Nacional de la Plata, CONICET, Museo de La Plata, Paseo del Bosque s/n, La Plata, Buenos Aires, Argentina*

*Corresponding author. Email: milen.marinov@mpi.govt.nz

‡These authors conceived the paper and contributed equally to the study.

- ⁵Muséum d'histoire naturelle, Département de Génétique et Evolution, Université de Genève, Geneva, Switzerland
- ⁶Biosecurity Surveillance & Incursion Investigation Plant Health Team, Ministry for Primary Industries, Christchurch, New Zealand
- ⁷Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil
- ⁸School of GeoSciences, University of Edinburgh, Edinburgh, Scotland, UK
- ⁹Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum National d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France
- ¹⁰Department of Biological Sciences, Indian Institute of Science Education and Research (IISER), Mohali, Punjab, India
- ¹¹Universidade do Vale do Rio dos Sinos (UNISINOS), São Leopoldo, Brazil
- ¹²Departamento de Entomología, Museo de Historia Natural de la Universidad Nacional Mayor de San Marcos, Lima, Peru
- ¹³Museo Regionale di Scienze Naturali, Torino, Italy
- ¹⁴Laboratório de Herpetologia, Departamento de Biodiversidade and Centro de Aquicultura (CAUNESP), Universidade Estadual Paulista-UNESP, Rio Claro, São Paulo, Brazil
- ¹⁵Department of Biological Sciences, National University of Singapore, Singapore
- ¹⁶Centro de Investigação em Biodiversidade e Recursos Genéticos - Rede de Investigação em Biodiversidade e Biologia Evolutiva, Universidade do Porto, Campus de Vairão, Vairão, Portugal
- ¹⁷Instituto de Biología Subtropical (CONICET-UNaM), Posadas, Misiones, Argentina
- ¹⁸Instituto Nacional da Biodiversidade e Áreas de Conservação, Luanda, Angola
- ¹⁹Federated Department of Biological Sciences, New Jersey Institute of Technology, Newark, New Jersey, USA
- ²⁰Instituto de Diversidad y Evolución Austral (IDEAus-CONICET), Puerto Madryn, Chubut, Argentina
- ²¹Laboratory of Soft Technologies, Valorization, Physico-Chemistry of Biological Materials and Biodiversity, Faculty of Science, University M'hamed Bougara of Boumerdes, Boumerdes, Algeria
- ²²Department of Biology, FFCLRP/ Universidade de São Paulo, Ribeirão Preto, Brazil
- ²³Section of Herpetology, Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany
- ²⁴691-8th Ave SE, Salmon Arm, British Columbia, V1E 2C2, Canada
- ²⁵Museu de Zoologia, Universidade de São Paulo, São Paulo, Brazil
- ²⁶Research and Collections, North Carolina Museum of Natural Sciences, Raleigh, North Carolina, USA
- ²⁷Department of Biology and Monte L. Bean Life Science Museum, Brigham Young University, Provo UT, USA
- ²⁸Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa
- ²⁹University of Tennessee at Chattanooga, Chattanooga, Tennessee, USA
- ³⁰The Linnean Society of London, Burlington House, Piccadilly, London, UK; Royal Botanic Gardens, Kew, Richmond, UK; Department of Environment and Agriculture, Curtin University, Perth, Australia
- ³¹División Plantas Vasculares, Universidad Nacional de la Plata, Museo de La Plata, Paseo del Bosque s/n, La Plata, Buenos Aires, Argentina
- ³²Instituto de Ciencias Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile
- ³³Marine and Environmental Sciences Centre, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal
- ³⁴Division of Invertebrate Zoology, American Museum of Natural History, New York, NY, USA
- ³⁵Pontifícia Universidade Católica do Rio Grande do Sul, Museu de Ciências e Tecnologia, Porto Alegre, Brazil
- ³⁶Estrada da Tundavala, Lubango, Angola
- ³⁷División Herpetología, Museo Argentino de Ciencias Naturales, 'Bernardino Rivadavia' – CONICET, Buenos Aires, Argentina
- ³⁸Department of Biological and Environmental Sciences, University of Gothenburg, Gothenburg, Sweden
- ³⁹Gantz Family Collection Center, Field Museum of Natural History, Chicago, USA
- ⁴⁰Società Italiana per la Storia della Fauna, Viale Liegi, Rome, Italy
- ⁴¹Herbário do Lubango, ISCED-Huíla, Rua Sarmento Rodrigues S/N, C.P. 230, Lubango, Angola

- ⁴²*Undergraduate Biology Program & Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, Kansas, USA*
- ⁴³*Department of Biological Sciences, University of Texas at El Paso, El Paso, Texas, USA*
- ⁴⁴*Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Mexico City, Mexico*
- ⁴⁵*Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu, China*
- ⁴⁶*Research Center for Biology, Indonesian Institute of Sciences, Jl. Raya Jakarta Bogor, Bogor, Indonesia*
- ⁴⁷*Departamento Invertebrados, Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, Brazil*
- ⁴⁸*Universidade Federal do Tocantins, Porto Nacional, Brazil*
- ⁴⁹*Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, Colombia*
- ⁵⁰*Institut Agro, Univ Angers, INRAE, IRHS, SFR QUASAV, Angers, France*
- ⁵¹*Museu Nacional de História Natural e da Ciência da Universidade de Lisboa, Lisboa, Portugal*
- ⁵²*Entomology Research Collection, Bio-Protection Research Centre, Lincoln University, Lincoln, New Zealand*
- ⁵³*Royal Saskatchewan Museum, Regina, Saskatchewan, Canada*
- ⁵⁴*Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil*
- ⁵⁵*Graduate School of Human and Environmental Sciences, Kyoto University, Kyoto, Japan*
- ⁵⁶*Tropical Herpetology Laboratory, Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Ilhéus, Brazil*
- ⁵⁷*Institute of Natural and Environmental Sciences, University of Hyogo, and Museum of Nature and Human Activities, Hyogo, Sanda, Japan*
- ⁵⁸*Centre for Functional Ecology. Science for People and the Planet, University of Coimbra, Portugal*
- ⁵⁹*Royal Belgian Institute of Natural Sciences, Brussels, Belgium*
- ⁶⁰*Laboratorio de Genética Evolutiva 'Claudio J. Bidau', Instituto de Biología Subtropical (IBS, CONICET), Universidad Nacional de Misiones (UNaM), Posadas, Misiones, Argentina*
- ⁶¹*Departamento de Biologia, Faculdade de Ciências da Universidade Agostinho Neto, Luanda, Angola*
- ⁶²*Department of Zoology, Faculty of Science, Charles University, Prague, Czech Republic*
- ⁶³*Department of Agroecology, Aarhus University, Denmark*
- ⁶⁴*Museum für Naturkunde – Leibniz Institute for Evolution and Biodiversity Science, Berlin, Germany*
- ⁶⁵*Instituto de Ciencias Marinas y Limnológicas, Facultad de Ciencias, Universidad Austral de Chile, Valdivia, Chile*
- ⁶⁶*Agrotechnology Study Program. Banau Tertiary Institute of Agricultural Enterprise, West Halmahera, Indonesia*
- ⁶⁷*Departament de Botànica i Geologia, Facultat de Ciències Biològiques, Universitat de València, Burjassot, València, Spain*
- ⁶⁸*Department of Plant Protection, Faculty of Agriculture, Suez Canal University, Ismailia, Egypt*
- ⁶⁹*Departamento de Biologia, Faculdade de Ciências da Universidade do Porto, Porto, Portugal*
- ⁷⁰*Department of Palaeontology and Historical Geology, Senckenberg Research Institute, Frankfurt am Main, Germany*
- ⁷¹*Grutas da Moeda e Fátima Lda., São Mamede, Batalha, Portugal*
- ⁷²*National Museum of Natural History, Bulgarian Academy of Sciences, Sofia, Bulgaria*
- ⁷³*Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia', Ciudad Autónoma de Buenos Aires, Argentina*
- ⁷⁴*Institut de Recherche pour le Développement (IRD), UMR MIVEGEC, Dakar, Senegal*
- ⁷⁵*Departamento de Biologia, Faculdade de Ciências da Universidade Agostinho Neto, Luanda, Angola*
- ⁷⁶*CSIR-Centre for Cellular and Molecular Biology, Laboratory for the Conservation of Endangered Species, Attapur, Hyderabad, India*
- ⁷⁷*Department of Pest-management and Conservation, Lincoln University, Lincoln, New Zealand*
- ⁷⁸*Society for Southeast Asian Herpetology, Heidelberg, Germany*
- ⁷⁹*Allwetterzoo, Münster, Germany*
- ⁸⁰*Hessisches Landesmuseum Darmstadt, Darmstadt, Germany*
- ⁸¹*American Museum of Natural History, New York, NY, USA*

⁸²*Senckenberg Forschungsstation Grube Messel, Senckenberg Forschungsinstitut und Naturmuseum Frankfurt am Main, Germany*

⁸³*Laboratory of Herpetology, Zoology Section, Department of Biology, Centre of Research in Natural Sciences at Lwiro, South Kivu Province, Democratic Republic of the Congo*

Received 18 June 2021; revised 22 July 2021; accepted for publication 25 July 2021

For almost 30 years, there have been active discussions about the taxonomic impediment and the challenge this represents to address the current human-induced biodiversity crisis. From the start (*Systematics Agenda 2000, 1994*), the term ‘taxonomic impediment’ has been ambiguous, designating both the insufficiency and inadequacy of the resources put to the service of taxonomy (the taxonomic impediment *sensu stricto*) and its main consequence, the wide discrepancy between the reality of specific biodiversity and our knowledge of it (the taxonomic gap; *Dubois, 2010; Raposo et al., 2020*). The total number of species on our planet is unknown, and its various estimates (using different methods) are widely divergent, but consensus exists that we are far from having inventoried half, and most likely one-tenth, of the species still present on earth today (*González-Oreja, 2008*).

Meanwhile, the biodiversity crisis has developed, so that it is now doubtless that a large part of these species will become extinct in the current and coming decades due to anthropogenic actions. This should put taxonomic urgency (*Wheeler et al., 2004; Dubois, 2010*) at the top of the priority list for biological sciences in our ‘century of extinctions’ (*Dubois, 2003*). It is clear that, despite its efforts, conservation biology will not significantly reduce the rate of extinction, even for vertebrates on which most attention is concentrated (*Hoffmann et al., 2010*). The main duty of scientists regarding this situation should be to reduce, as much as possible, the taxonomic gap and to store specimens and tissues that will testify to the vanishing biodiversity of our planet. Once safely stored, research on this material will be possible, even if the species have fallen to extinction in the interim.

Taxonomy is an undervalued biological discipline. This is mainly due to several misconceptions about taxonomy that are widespread in the scientific community. Among these are: (1) the idea that, because taxonomy uses ‘type specimens’, the discipline is typological and non-evolutionary; (2) that taxonomists only study morphology; or (3) that their work is ancillary, merely providing identification of specimens used in other biological studies, such as molecular phylogeny, ecology or conservation biology.

So-called ‘type specimens’ are in fact just vouchers that serve as ‘name-bearers’, allowing objective allocation of names to taxa and are not meant as being ‘typical’ of the taxon (*Simpson, 1940*). Taxonomy, as with every other science, is based on explicit concepts and on the formulation and testing of hypotheses. It requires an integrative approach where scientists benefit by working collaboratively to synthesize the results of data derived from morphology, behaviour, bioacoustics, ecology, geography, molecules, chemistry, history and other diverse fields of inquiry. In this sense it is perhaps the most integrative of all biological disciplines (*de Carvalho et al., 2008*). The identification of specimens is a (useful) practical outcome of taxonomy, but not the core of the discipline, which consists of taxonomic revisions in which the circumscriptions and relationships of taxa are (re)-evaluated (*Löbl, 2018*). Taxonomy is indeed a descriptive science that tests explicit hypotheses, in the same fashion as stellar science and planetary probes represent nothing more than high-tech descriptive sciences (*Grimaldi & Engel, 2007*). Of course, most great finds in science, from gravity to evolution, stem from description of observations.

The source of the ‘taxonomic impediment’ is unclear, but is perceived by some as meaning that the pace of taxonomic work is too slow and laborious, and restricted to the painstaking study of morpho-anatomy, thus insufficient to meet the challenge of documenting biodiversity before it is driven to extinction by anthropogenic alteration and destruction of our ecosystems. Accordingly, various attempts to ‘innovate and speed up’ the taxonomic process have emerged. Some simplistic proposals, mostly based on technological approaches, have been presented as ‘revolutionary’, for example, replacing descriptions and hypotheses of species with a DNA barcode (wrongly dubbed a ‘diagnosis’; *Sharkey et al., 2021*), or replacing name-bearing specimens held in public curated collections with photographs of species live in the field (*Cerriaco et al., 2016*).

The difficulty in appropriately understanding and addressing the taxonomic crisis results from its multifactorial causes. Among them, the following three are particularly important (*de Carvalho et al., 2005*;

The Buffon Symposium, 2008; Boero, 2010; Wheeler, 2014; Prathapan *et al.*, 2018; Britz *et al.*, 2020; Dubois *et al.*, 2021): (1) the lack of recognition of taxonomy as a robust science by peers and policy-makers; (2) the considerable lack of well-trained professionals, having both solid theoretical and practical backgrounds; (3) the many legal, administrative and ideological barriers that exist regarding the collection and storing of specimens, without which no proper scientific taxonomy can exist.

Many of the problems of taxonomy stem from the dismissive attitude of the scientific community toward this scientific discipline, its unwillingness to appropriately support taxonomic work, a concomitant elimination of academic positions and a growing suspicion towards fieldwork and specimen collection. The search for ‘miracle solutions’ to taxonomy is mostly based on technology rather than manpower. Technological innovations may bring useful tools aiding taxonomic work, but will never replace it. If the population of taxonomists was growing, and if they were given the financial and practical means to carry out well-organized field surveys and collection campaigns worldwide (Dubois *et al.*, 2021), then the taxonomic impediment could rapidly erode away.

The poor image for taxonomy among biological sciences, its concomitant devaluation of taxonomic publications (taxonomic treatments, monographs and new species publications are rarely cited outside taxonomy, even if they are used by other disciplines), and shortage of professional positions and research grants, have played a devastating role in the development and recruitment for the discipline.

There is a global lack of academic education in taxonomy and of properly trained taxonomists, that is, scientists who distinguish and classify taxa according to explicit concepts and data-based hypotheses, and name them respecting nomenclatural rules and checking type specimens. The theoretical weaknesses of some of the simplistic approaches mentioned above testify to the ignorance of their supporters of the basic tenets of scientific taxonomy (Amorim *et al.*, 2016; Santos *et al.*, 2016). Regardless of technical problems related to the naming of the ‘new taxa’ erected by ‘revolutionary’ protocols (Bauer *et al.*, 2011), such proposals are scientifically questionable.

The case of DNA barcoding is emblematic in this respect. This technique may be useful to identify specimens, that is, to allocate them to species already recognized by proper taxonomic work or to associate otherwise challenging polymorphic or cryptic taxa. As such, it may be useful in ecology, conservation biology, environmental monitoring and other domains, but it is inappropriate as the sole basis for taxonomic work, which consists in delimitating species and ascertaining their relationships. The use of arbitrary

‘threshold values’ as measured by this methodology to ‘distinguish species’ is nothing but a recent avatar of phenetic taxonomic methods that were dismissed long before the onset of DNA sequencing.

Such approaches actually worsen the challenge by addressing a red herring lurking in the notion of the taxonomic impediment. Moreover, they are based on the assumption that the goal of taxonomy is the mere placing of a name in a taxonomic hierarchy rather than acknowledging that the number, although interesting, lacks practical value in addressing the biodiversity challenges of today (de Carvalho *et al.*, 2014). The goal is not to build mere lists of taxa, just as listing diseases or chemical elements are not the goals of medicine or chemistry, but such lists are integral to all of these disciplines. However, names divorced from suitable breadths of adequate information confuse rather than advance our agendas. Instead, we need means to efficiently, accurately and quickly (but not compromising the former actions) establish the existence of discrete evolutionary units called species, which will allow for their later identification in the lab and field, as well as an understanding of how those species interact with their environment. A cavalierly produced list of species with names dissociated from proper scientific species concepts is no more valuable than knowing the number of different pills in a jar, but failing to know which are stimulants, depressants, painkillers or hallucinogens.

Aside from the aforementioned considerable shortage of brains and hands, a major factor in the taxonomic crisis results from the difficulties encountered by taxonomists to study and collect specimens in the field, store them safely in well-curated collections and make them available to colleagues for study. For a significant increase of our knowledge of biodiversity, collecting new specimens in the field will remain the rate-limiting step (May, 2004). New species are not found in the computers and sequencers of the big cities of the ‘North’, but in the natural and anthropized habitats of the entire planet and, especially, in the more biodiverse ‘South’ (Dubois *et al.*, 2021). Molecular phylogenetics, taxonomic databases and ‘big data’ derived from existing taxonomic information are, indeed, useful tools, but they do not increase our current knowledge of the species that have not yet been collected or studied. In our century of extinctions, immediate effort must be dedicated to collaborative fieldwork and collections, while there is still time.

Organic evolution is not teleological. It results from a variable combination of ‘chance and necessity’ (Monod, 1970), that is, of genetic variation and natural selection, and as such is not deterministic and predictable. The only way to know it in-depth is to study the organisms themselves, not merely ‘models’ derived from a superficial knowledge of small subsamples.

Efforts to establish new taxa based on the structure of a fragment of one particular gene, or to replace biological vouchers with photographs, actually exacerbate the taxonomic impediment. As in all sciences, there is a need for best practices. In order to attend to the current crises, we need more than ever an adherence to the best standards for producing taxonomic science and communicating it using the proper technical language of scientific nomenclature. This includes a synthesis of extensive data sources to support taxon circumscriptions and hypotheses, and suitable vouchers for their later verification and scientific replication. Specimens are needed so that future generations have a record of life at present, sometimes the sole data we shall ever have for species if it is soon to be lost. Future generations may use these specimens to verify our hypotheses in the face of changing concepts of species and gather data that today are undreamt of – in the same way we today may extract DNA or employ CT-scan data of Linnean or Darwinian specimens to gain insights that those historical collectors could never have imagined themselves.

Instead of relying solely on barcodes and/or photographs, we need intentional and mindful syntheses of molecular, morphological and other data to aid the progress of our knowledge of the vanishing biodiversity. Recent ‘revolutionary protocols’, if adopted by the community, would make the toil of every taxonomist far greater (Meier *et al.*, *In press*). Taxonomists would be faced with a tremendous hurdle toward actually identifying thousands of specimens that may pass before them and may be crucial for ecological, conservation, agricultural or climate-change studies. Every specimen would require partial or complete destruction, depending on the size of the sampled organisms, in order to match the DNA diagnoses, transforming an afternoon’s worth of study into days or weeks. Rather than moving on to tackle the next taxonomic challenge, our diminished force of taxonomists would be bogged down in attempts to place specimens into seemingly known species, with the potential for greater errors if any should attempt to rely solely on photographs, which in many cases fail to properly allow for distinguishing between closely similar or cryptic taxa.

Best practices in taxonomy are critical for addressing the current biodiversity crisis. Innovation is good, but it should improve on an existing situation rather than exacerbate difficulties or worse, revert to 18th- or 19th-century one-character (barcode or single gene) taxonomic approaches. Taxonomy needs to be rebranded to the scientific community as a modern, active and important discipline, and taxonomists need to come together to demand sufficient funding, training and employment for taxonomic researchers.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

- Amorim DS, Santos CM, Krell FT, Dubois A, Nihei SS, Oliveira OM, Pont A, Song H, Verdade VK, Fachin DA, Klassa B, Lamas CJ, Oliveira SS, Carvalho CJ, Mello-Patiu CA, Hajdu E, Couri MS, Silva VC, Capellari RS, Falaschi RL, Feitosa RM, Prendini L, Pombal JP, Fernández F, Rocha RM, Lattke JE, Caramaschi U, Duarte M, Marques AC, Reis RE, Kurina O, Takiya DM, Tavares M, Fernandes DS, Franco FL, Cuezso F, Paulson D, Guénard B, Schlick-Steiner BC, Arthofer W, Steiner FM, Fisher BL, Johnson RA, Delsinne TD, Donoso DA, Mulieri PR, Patitucci LD, Carpenter JM, Herman L, Grimaldi D. 2016. Timeless standards for species delimitation. *Zootaxa* **4137**: 121–128.
- Bauer AM, Parham JF, Brown RM, Stuart BL, Grismer L, Papenfuss TJ, Böhme W, Savage JM, Carranza S, Grismer JL, Wagner P, Schmitz A, Ananjeva NB, Inger RF. 2011. Availability of new Bayesian-delimited gecko names and the importance of character-based species descriptions. *Proceedings of the Royal Society, B, Biological Sciences* **278**: 490–492; discussion 493.
- Boero F. 2010. The study of species in the era of biodiversity: a tale of stupidity. *Diversity* **16**: 115–126.
- Britz R, Hundsdoerfer A, Fritz U. 2020. Funding, training, permits – the three big challenges of taxonomy. *Megataxa* **1**: 49–52.
- de Carvalho MR, Bockmann FA, Amorim DS, de Vivo M, de Toledo-Piza M, Menezes NA, de Figueiredo JL, Castro RM, Gill AC, McEachran JD, Compagno LJ, Schelly RC, Britz R, Lundberg JG, Vari RP, Nelson G. 2005. Revisiting the taxonomic impediment. *Science (New York, N.Y.)* **307**: 353.
- de Carvalho MR, Bockmann FA, Amorim DS, Brandão CRF. 2008. Systematics must embrace comparative biology and evolution, not speed and automation. *Evolutionary Biology* **35**: 150–157.
- de Carvalho MR, Ebach MC, Williams DM, Nihei SS, Trefaut Rodrigues M, Grant T, Silveira LF, Zaher H, Gill AC, Schelly RC, Sparks JS, Bockmann FA, Séret B, Ho H-C, Grande L, Rieppel O, Dubois A, Ohler A, Faivovich J, Assis LCS, Wheeler QD, Goldstein PZ, de Almeida EAB, Valdecasas AG, Nelson G. 2014. Does counting species count as taxonomy? On misrepresenting systematics, yet again. *Cladistics* **30**: 322–329.
- Ceríaco LM, Gutiérrez EE, Dubois A, Carr M; + 490 signatories. 2016. Photography-based taxonomy is inadequate, unnecessary, and potentially harmful for biological sciences. *Zootaxa* **4196**: 435–445.
- Dubois A. 2003. The relationships between taxonomy and conservation biology in the century of extinctions. *Comptes Rendus Biologies* **326** Suppl 1: S9–S21.

- Dubois A. 2010.** Zoological nomenclature in the century of extinctions: priority vs. usage. *Organisms, Diversity & Evolution* **10**: 259–274.
- Dubois A, Ohler A, Pyron RA. 2021.** New concepts and methods for phylogenetic taxonomy and nomenclature in zoology, exemplified by a new ranked cladonomy of recent amphibians (Lissamphibia). *Megataxa* **5**: 1–738.
- González-Oreja JA. 2008.** The encyclopedia of life vs. the brochure of life: exploring the relationships between the extinction of species and the inventory of life on Earth. *Zootaxa* **1965**: 61–68.
- Grimaldi DA, Engel MS. 2007.** Why descriptive science still matters. *BioScience* **57**: 646–647.
- Hoffmann M, Hilton-Taylor C, Angulo A, Böhm M, Brooks TM, Butchart SH, Carpenter KE, Chanson J, Collen B, Cox NA, Darwall WR. 2010.** The impact of conservation on the status of the world's vertebrates. *Science* **330**:1503–1509.
- Löbl I. 2018.** Assessing biotic diversity: the glorious past, present, and the uncertain future. *Bulletin of the Entomological Society of Malta* **10**: 5–15.
- May RM. 2004.** Tomorrow's taxonomy: collecting new species in the field will remain the rate-limiting step. *Philosophical Transactions of the Royal Society of London, Series B, Biological sciences* **359**: 733–734.
- Meier R, Blaimer B, Buenaventura E, Hartop E, von Rintelen T, Srivathsan A, Yeo D. In press.** A re-analysis of the data in Sharkey *et al.*'s (2021) minimalist revision reveals that BINs do not deserve names, but BOLD Systems needs a stronger commitment to open science. *Cladistics*.
- Monod J. 1970.** *Le hasard et la nécessité. Essai sur la philosophie naturelle de la biologie moderne*. Paris: Seuil, 1–213.
- Prathapan KD, Pethiyagoda R, Bawa KS, Raven PH, Rajan PD; + 172 co-signatories from 35 countries. 2018.** When the cure kills – CBD limits biodiversity research. *Science (New York, N.Y.)* **360**: 1405–1406.
- Raposo MA, Kirwan GM, Lourenço ACC, Sobral G, Bockmann FA, Stopiglia R. 2020.** On the notions of taxonomic 'impediment', 'gap', 'inflation' and 'anarchy', and their effects on the field of conservation. *Systematics & Biodiversity* **19**: 296–311.
- Santos CMD, Amorim DS, Klassa B, Fachin DA, Nihei SS, Carvalho CJB de, Falaschi RL, Mello-Pitau CA, Couri MS, Oliveira SS, Silva VC, Ribeiro GC, Capellari RS, Lamas CJE. 2016.** On typeless species and the perils of fast taxonomy. *Systematic Entomology* **41**: 511–515.
- Sharkey MJ, Janzen DH, Hallwachs W, Chapman EG, Smith MA, Dapkey T, Brown A, Ratnasingham S, Naik S, Manjunath R, Perez K, Milton M, Hebert P, Shaw SR, Kittel RN, Alma Solis M, Metz MA, Goldstein PZ, Brown JW, Quicke DLJ, Van Achterberg C, Brown BV, Burns JM. 2021.** Minimalist revision and description of 403 new species in 11 subfamilies of Costa Rican braconid parasitoid wasps, including host records for 219 species. *ZooKeys* **1013**: 1–665.
- Simpson GG. 1940.** Types in modern taxonomy. *American Journal of Science* **238**: 413–431.
- Systematics Agenda 2000. 1994.** *Charting the biosphere: a global initiative to discover, describe and classify the world's species. Technical report*. New York: American Museum of Natural History, American Society of Plant Taxonomy, Society of Systematic Biologists and the Willi Hennig Society, 1–34.
- The Buffon Symposium. 2008.** *The Buffon Declaration. Natural history institutions and the environmental crisis. Concluding message from the Buffon Symposium – October 18th and 19th, 2007*. Paris: Muséum national d'Histoire Naturelle. Available from: http://www.mnhn.fr/museum/front/medias/pdf/13696_The_Buffon_Declaration_CBDmay08_Eng.pdf. Accessed 24 October 2007.
- Wheeler Q. 2014.** Are reports of the death of taxonomy an exaggeration? *New Phytologist* **201**: 370–371.
- Wheeler QD, Raven PH, Wilson EO. 2004.** Taxonomy: impediment or expedient? *Science (New York, N.Y.)* **303**: 285.